

BEREZIN TRANSFORM IN POLYNOMIAL BERGMAN SPACES

YACIN AMEUR, HÅKAN HEDENMALM, AND NIKOLAI MAKAROV

ABSTRACT. We consider fairly general weight functions $Q : \mathbb{C} \rightarrow \mathbb{R}$, and let $K_{m,n}$ denote the reproducing kernel for the space $H_{m,n}$ of analytic polynomials u of degree at most $n-1$ of finite norm $\|u\|_{mQ}^2 = \int_{\mathbb{C}} |u(z)|^2 e^{-mQ(z)} dA(z)$, dA denoting suitably normalized area measure in $\mathbb{C}$. For a continuous bounded function f on $\mathbb{C}$, we consider its (polynomial) Berezin transform

$$\mathfrak{B}_{m,n}f(z) = \int_{\mathbb{C}} f(\zeta) dB_{m,n}^{(z)}(\zeta) \quad \text{where} \quad dB_{m,n}^{(z)}(\zeta) = \frac{|K_{m,n}(z, \zeta)|^2}{K_{m,n}(z, z)} e^{-mQ(\zeta)} dA(\zeta).$$

Let $X = \{\Delta Q > 0\}$. For a parameter $\tau > 0$ we prove that there exists a compact subset S_τ of $\mathbb{C}$ such that

$$(0.1) \quad \mathfrak{B}_{m,n}f(z) \rightarrow f(z) \quad \text{as } m \rightarrow \infty \text{ and } n - m\tau \rightarrow 0,$$

for all continuous bounded f if z is in the interior of $S_\tau \cap X$. Equivalently, the measures $B_{m,n}^{(z)}$ converge to the Dirac measure at z . The set S_τ is the coincidence set for an associated obstacle problem.

We also prove that the convergence in (0.1) is *Gaussian* when z is in the interior of $S_\tau \cap X$, in the sense that with $F(\zeta) = f(\sqrt{m\Delta Q(\zeta)}(\zeta - z))$, we have

$$\mathfrak{B}_{m,n}F(z) \rightarrow \int_{\mathbb{C}} f(\zeta) dP(\zeta),$$

$dP(\zeta) = e^{-|\zeta|^2} dA(\zeta)$ denoting the standard Gaussian.

In the "model case" $Q(z) = |z|^2$, S_τ is the closed disk with centre 0 and radius $\sqrt{\tau}$. We prove that if z is fixed with $|z| > \sqrt{\tau}$, the corresponding measures $B_{m,n}^{(z)}$ converge to harmonic measure for z relative to the domain $\mathbb{C}^* \setminus S_\tau$, $\mathbb{C}^*$ denoting the extended plane.

Our auxiliary results include L^2 estimates for the $\bar{\partial}$ -equation $\bar{\partial}u = f$ when f is a suitable test function and the solution u is restricted by a growth constraint near ∞ .

Our results have applications e.g. to the study of weighted distributions of eigenvalues of random normal matrices. In the companion paper [1] we consider such applications, e.g. a proof of Gaussian field convergence for fluctuations of linear statistics of eigenvalues of random normal matrices from the ensemble associated with Q .

1. INTRODUCTION

1.1. General introduction to Berezin quantization.¹

In a version of quantum theory, a Bargmann-Fock type space of polynomials plays the role of the quantized system, while the corresponding weighted L^2 space is the classical analogue. It is therefore natural to study the asymptotics of the quantized system as we approach the semiclassical limit. A particularly useful object is the *reproducing kernel* of the Bargmann-Fock type space. To make matters more concrete, let μ be a finite positive Borel measure on $\mathbb{C}$, and

Research supported by the Göran Gustafsson Foundation. The third author is supported by N.S.F. Grant No. 0201893.

¹In this version we have but slightly modified the formulation of Theorem 2.8 and made some minor corrections.

$L^2(\mathbb{C}; \mu)$ the usual L^2 space with inner product

$$\langle f, g \rangle_{L^2(\mathbb{C}; \mu)} = \int_{\mathbb{C}} f(z) \overline{g(z)} d\mu(z).$$

The subspace of $L^2(\mathbb{C}; \mu)$ of entire functions is the Bergman space $A^2(\mathbb{C}; \mu)$.

We assume that the support of μ has infinitely many points so that any given polynomial corresponds to a unique element of $L^2(\mathbb{C}; \mu)$, and write

$$J_\mu = \sup \left\{ j \in \mathbb{Z}; \int_{\mathbb{C}} |z|^{2(j-1)} d\mu(z) < \infty \right\}.$$

Since μ is finite, we are ensured that $1 \leq J_\mu \leq \infty$.

Let $\mathcal{P}_n$ be the set of analytic polynomials of degree at most $n-1$, and write

$$A_{\mu,n}^2 = L^2(\mathbb{C}; \mu) \cap \mathcal{P}_n \subset L^2(\mathbb{C}; \mu).$$

Putting $n' = \min\{n, J_\mu\}$, it is thus seen that $A_{\mu,n}^2$ equals $\mathcal{P}_{n'}$ in the sense of sets, with the norm inherited from $L^2(\mathbb{C}; \mu)$. Hence $A_{\mu,n}^2 = A_{\mu,n'}^2$ for all n , and the reader who so desires can without loss of generality assume that $n = n'$ in the following.

Let $e_1, \dots, e_{n'}$ be an orthonormal basis for $A_{\mu,n}^2$. The reproducing kernel $K_{\mu,n}$ for the space $A_{\mu,n}^2$ is the function

$$K_{\mu,n}(z, w) = \sum_{j=1}^{n'} e_j(z) \overline{e_j(w)}.$$

Then $K_{\mu,n}$ is independent of the choice of an orthonormal basis for $A_{\mu,n}^2$, and it is characterized by the properties that $z \mapsto K_{\mu,n}(z, z_0)$ is in $A_{\mu,n}^2$ and

$$(1.1) \quad u(z_0) = \langle u, K_{\mu,n}(\cdot, z_0) \rangle_{L^2(\mathbb{C}; \mu)} = \int_{\mathbb{C}} u(z) \overline{K_{\mu,n}(z, z_0)} d\mu(z), \quad u \in A_{\mu,n}^2, \quad z_0 \in \mathbb{C}.$$

For a given complex number z_0 we now consider the measure

$$(1.2) \quad dB_{\mu,n}^{\langle z_0 \rangle}(z) = \frac{|K_{\mu,n}(z, z_0)|^2}{K_{\mu,n}(z_0, z_0)} d\mu(z),$$

which we may call the *Berezin measure* associated with μ , n and z_0 . The reproducing property (1.1) applied to $u = K_{\mu,n}(\cdot, z_0)$ implies that $B_{\mu,n}^{\langle z_0 \rangle}$ is a probability measure. In classical physics, $B_{\mu,n}^{\langle z_0 \rangle}$ corresponds to a point mass at z_0 , so our interest focuses on how closely this measure approximates the point mass.

1.2. Weights. By a *weight* we mean a measurable function $\phi : \mathbb{C} \rightarrow \mathbb{R}$ such that the measure

$$d\mu_\phi(z) = e^{-\phi(z)} dA(z),$$

is a finite measure on $\mathbb{C}$, where $dA(z) = dx dy / \pi$ with $z = x + iy$.

We write L_ϕ^2 for the space $L^2(\mathbb{C}; \mu_\phi)$ and A_ϕ^2 for $A^2(\mathbb{C}; \mu_\phi)$ and the norm of an element $u \in L_\phi^2$ will be denoted by

$$\|u\|_\phi^2 = \int_{\mathbb{C}} |u|^2 e^{-\phi} dA.$$

For a positive integer n , we frequently write

$$(1.3) \quad L_{\phi,n}^2 = \left\{ u \in L_\phi^2; u(z) = O(|z|^{n-1}) \text{ when } |z| \rightarrow \infty \right\}$$

and

$$(1.4) \quad A_{\phi,n}^2 = L_{\phi,n}^2 \cap A_\phi^2 = L_{\phi,n}^2 \cap \mathcal{P}_n.$$

Observe that $L_{\phi,n}^2$ usually is non-closed in L_ϕ^2 , whereas the finite-dimensional $A_{\phi,n}^2$ is closed in L_ϕ^2 .

1.3. The weights considered. Let $Q : \mathbb{C} \rightarrow \mathbb{R}$ be a given measurable function, which satisfies a growth condition of the form

$$(1.5) \quad Q(z) \geq \rho \log |z|^2, \quad |z| \geq C,$$

for some positive numbers ρ and C .

For a positive number m , we now consider weights ϕ_m of the form $\phi_m = mQ$. By abuse of notation, we will in this context sometimes refer to Q as the weight. To simplify notation, we set

$$H_{m,n} = A_{mQ,n}^2 = L_{mQ}^2 \cap \mathcal{P}_n$$

and denote by $K_{m,n}$ the reproducing kernel for $H_{m,n}$. For a given z_0 , the corresponding $B_{m,n}^{(z_0)}$ is defined accordingly, cf. (1.2).

We think of Q as being fixed while the parameters m and n vary, and also fix a number τ satisfying $0 < \tau < \rho$. To avoid bulky notation, it is customary to reduce the number of parameters to one, by regarding $n = n(m)$ as a function of $m > 0$. We will adopt this convention and study the behaviour of the measures $B_{m,n}^{(z_0)}$ when $m \rightarrow \infty$ and $n - m\tau \rightarrow 0$.

Adding a real constant to Q means that the inner product in $H_{m,n}$ is only changed by a positive multiplicative constant, and $K_{m,n}$ gets multiplied by the inverse of that number. Hence the corresponding Berezin measures are unchanged, and we can for example w.l.o.g. assume that $Q \geq 1$ on $\mathbb{C}$ when this is convenient.

Remark 1.1. Replacing Q by $\tau^{-1}Q$ one can assume that $\tau = 1$. The most important case (e.g. in random matrix theory) occurs when $\tau = 1$ and $m = n$. However, we shall take on the slightly more general approach (with three parameters m, n and τ , instead of just n) in this note.

1.4. A word on notation. For real x , we write $\lfloor x \rfloor$ for the largest integer which is strictly smaller than x . We write frequently

$$\partial_z = \frac{\partial}{\partial z} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \bar{\partial}_z = \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right), \quad z = x + iy,$$

and use $\Delta = \Delta_z$ to denote the normalized Laplacian

$$\Delta_z = \partial_z \bar{\partial}_z = \frac{1}{4} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right).$$

We write $\mathbb{D}(z_0; r)$ for the open disk $\{z \in \mathbb{C}; |z - z_0| < r\}$, and we simplify the notation to $\mathbb{D}$ when $z_0 = 0$ and $r = 1$. The boundary of $\mathbb{D}(z_0, r)$ is denoted $\mathbb{T}(z_0, r)$. When S is a subset of $\mathbb{C}$, we write S° for the interior of S , and $\bar{S}$ for its closure; the support of a continuous function f is denoted by $\text{supp } f$. The symbol $A \subset B$ means that A is a subset of B , while $A \Subset B$ means that A is a compact subset of B . We write $\text{dist}(A, B)$ for the Euclidean distance between A and B . The symbol *a.e.* is short for "dA-almost everywhere", and *p.m.* is short for "probability measure". Finally, we use the shorthand L^2 to denote the unweighted space $L_0^2 = L^2(\mathbb{C}; dA)$.

1.5. Random normal matrices and the Coulomb gas. This paper is a slight reworking of a preprint from arxiv.org which was written in 2007 and early 2008. During the work, it became convenient to collect, in a separate paper, some estimates needed for the paper [1], which was written simultaneously. More precisely, we wanted to make clear a variant of some results on asymptotic expansions of (polynomial) Bergman kernels which were known in a several variables context ([6], [3]), as well as some uniform estimates in the off-diagonal case. The relevant results for our applications in [1] are primarily theorems 2.8 and 8.3. A discussion containing related estimates which hold near the boundary will appear in our later publication [2].

2. MAIN RESULTS

2.1. Berezin quantization. Fix a non-negative weight Q and two positive numbers τ and ρ , $\tau < \rho$, such that the growth condition (1.5) is satisfied, and form the measures

$$dB_{m,n}^{(z_0)} = B_{m,n}^{(z_0)} dA \quad \text{where} \quad B_{m,n}^{(z_0)}(z) = \frac{|K_{m,n}(z, z_0)|^2}{K_{m,n}(z_0, z_0)} e^{-mQ(z)}.$$

We shall refer to the function $B_{m,n}^{(z_0)}$ as the *Berezin kernel* associated with m, n and z_0 .

Let us now assume that $Q \in C^2(\mathbb{C})$. In the first instance, we ask for which z_0 we have convergence

$$B_{m,n}^{(z_0)} \rightarrow \delta_{z_0} \quad \text{as } m \rightarrow \infty \text{ and } n - m\tau \rightarrow 0,$$

in the sense of measures. In terms of the *Berezin transform*

$$\mathfrak{B}_{m,n} f(z_0) = \int_{\mathbb{C}} f(z) dB_{m,n}^{(z_0)}(z),$$

we are asking whether

$$(2.1) \quad \mathfrak{B}_{m,n} f(z_0) \rightarrow f(z_0) \quad \text{for all } f \in C_b(\mathbb{C}) \text{ as } m \rightarrow \infty \text{ and } n - m\tau \rightarrow 0.$$

Let X be the set of points where Q is strictly subharmonic,

$$X = \{\Delta Q > 0\}.$$

We shall find that there exists a compact set $\mathcal{S}_\tau$ such that (2.1) holds for all z_0 in the interior of $\mathcal{S}_\tau \cap X$, while (2.1) fails whenever $z_0 \notin \mathcal{S}_\tau$.

To define $\mathcal{S}_\tau$, we first need to introduce some notions from weighted potential theory, cf. [21] and [16]. It is here advantageous to slightly relax the regularity assumption on Q and assume that Q is in the class $C^{1,1}(\mathbb{C})$ consisting of all C^1 -smooth functions with Lipschitzian gradients. We will have frequent use of the simple fact that the distributional Laplacian ΔF of a function $F \in C^{1,1}(\mathbb{C})$ makes sense almost everywhere and is of class $L_{\text{loc}}^\infty(\mathbb{C})$.

Let SH_τ denote the set of all subharmonic functions $f : \mathbb{C} \rightarrow \mathbb{R}$ which satisfy a growth bound of the form

$$f(z) \leq \tau \log |z|^2 + O(1) \quad \text{as } z \rightarrow \infty.$$

The *equilibrium potential* corresponding to Q and the parameter τ is defined by

$$\widehat{Q}_\tau(z) = \sup \{f(z); f \in \text{SH}_\tau \text{ and } f \leq Q \text{ on } \mathbb{C}\}.$$

Clearly, $\widehat{Q}_\tau$ is subharmonic on $\mathbb{C}$. We now define $\mathcal{S}_\tau$ as the coincidence set

$$(2.2) \quad \mathcal{S}_\tau = \{Q = \widehat{Q}_\tau\}.$$

Evidently, $\mathcal{S}_\tau$ increases with τ , and that Q is subharmonic on $\mathcal{S}_\tau^\circ$.

We shall need the following lemma.

Lemma 2.1. *The function $\widehat{Q}_\tau$ is of class $C^{1,1}(\mathbb{C})$, and $\mathcal{S}_\tau$ is compact. Furthermore, $\widehat{Q}_\tau$ is harmonic in $\mathbb{C} \setminus \mathcal{S}_\tau$ and there is a number C such that $\widehat{Q}_\tau(z) \leq \tau \log_+ |z|^2 + C$ for all $z \in \mathbb{C}$.*

Proof. This is [[16], Prop. 4.2, p. 10] and [[21], Th. I.4.7, p. 54]. $\square$

It is easy to construct subharmonic minorants of critical growth; for C large enough, the function $z \mapsto \tau \log_+ (|z|^2 / C)$ is a minorant of Q of class SH_τ . It yields that

$$(2.3) \quad \widehat{Q}_\tau(z) = \tau \log |z|^2 + O(1) \quad \text{as } z \rightarrow \infty.$$

Proposition 2.2. *Let $Q \in C^{1,1}(\mathbb{C})$ and $z_0 \in \mathbb{C}$ an arbitrary point. Then $B_{m,n}^{(z_0)}(\Lambda) \rightarrow 1$ as $m \rightarrow \infty$ and $n \leq m\tau + 1$ for every open neighbourhood Λ of $\mathcal{S}_\tau$.*

Proof. See Sect. 3. □

It follows from Prop. 2.2 that $B_{m,n}^{(z_0)}(\mathbb{C} \setminus \Lambda) \rightarrow 0$. Hence if $z_0 \notin S_\tau$, then (2.1) fails in general, and

$$\mathfrak{B}_{m,n}f(z_0) \rightarrow 0 \quad \text{as } m \rightarrow \infty \quad \text{and } n \leq m\tau + 1,$$

for all $f \in C_b(\mathbb{C})$ such that $\text{supp } f \cap S_\tau = \emptyset$. The situation is entirely different when the point z_0 is in the interior of $S_\tau \cap X$.

Theorem 2.3. *Assume that $Q \in C^2(\mathbb{C})$ and let $z_0 \in S_\tau^\circ \cap X$. Then, for any $f \in C_b(\mathbb{C})$, and any real number $M \geq 0$, the measures $B_{m,n}^{(z_0)}$ converge to δ_{z_0} as $m \rightarrow \infty$ and $n \geq m\tau - M$.*

Proof. See Sect. 7. See also Rem. 2.5. □

2.2. A more elaborate estimate for the Berezin kernel. Th. 2.3 suggests that if z_0 is a given point in the interior of $S_\tau \cap X$, m is large, and $n \geq m\tau - 1$, then the density $B_{m,n}^{(z_0)}(z)$ should attain its maximum for z close to z_0 . The following theorem implies that this is the case, and gives a good control for $B_{m,n}^{(z_0)}$ in the critical case, when $n - m\tau \rightarrow 0$.

Theorem 2.4. *Assume that $Q \in C^2(\mathbb{C})$. Let K be a compact subset of $S_\tau^\circ \cap X$ and fix a non-negative number M . Put*

$$d_K = \text{dist}(K, \mathbb{C} \setminus (S_\tau \cap X)) \quad \text{and} \quad a_K = \inf\{\Delta Q(z); \text{dist}(z, K) \leq d_K/2\}.$$

Then there exists positive numbers C and ϵ such that

$$B_{m,n}^{(z_0)}(z) \leq Cme^{-\epsilon\sqrt{m}\min\{d_K, |z-z_0|\}}e^{-m(Q(z)-\widehat{Q}_\tau(z))}, \quad z_0 \in K, z \in \mathbb{C}, m\tau - M \leq n \leq m\tau + 1.$$

Here C depends only on K , M and τ , while ϵ only depends on M , a_K and τ .

Proof. See Sect. 8. □

A similar result was proved independently by Berman [4] after the completion of this note.

Remark 2.5. For fixed M and τ , the number ϵ can be chosen proportional to a_K . A related result with much more precise information on the dependence of C and ϵ on the various parameters in question is given in Th. 8.1 and the subsequent remark.

We also remark that our proof for Th. 2.4 is very different from that for Th. 2.3. Thus Th. 2.4 gives an alternative method for obtaining Th. 2.3 in the critical case when $n - m\tau \rightarrow 0$.

2.3. Gaussian convergence. Fix a point $z_0 \in X$. It will be convenient to introduce the *normalized* Berezin measure $\widehat{B}_{m,n}^{(z_0)}$ by

$$(2.4) \quad d\widehat{B}_{m,n}^{(z_0)} = \widehat{B}_{m,n}^{(z_0)} dA \quad \text{where} \quad \widehat{B}_{m,n}^{(z_0)}(z) = \frac{1}{m\Delta Q(z_0)} B_{m,n}^{(z_0)}\left(z_0 + \frac{z}{\sqrt{m\Delta Q(z_0)}}\right).$$

We denote the standard Gaussian p.m. on $\mathbb{C}$ by

$$dP(z) = e^{-|z|^2} dA(z).$$

We have the following CLT, which gives much more precise information than Th. 2.3. We will settle for stating it for C^∞ weights Q which are *real-analytic* in a neighbourhood of S_τ , but remark that, using well-known methods, our proof can be extended to cover the case of, say, C^∞ -smooth weights. We will give further details on possible generalizations later on.

Theorem 2.6. *Assume that Q is real-analytic in a neighbourhood of S_τ . Fix a compact subset $K \Subset S_\tau^\circ \cap X$, a point $z_0 \in K$, and a number $M \geq 0$. Then we have*

$$(2.5) \quad \int_{\mathbb{C}} \left| \widehat{B}_{m,n}^{(z_0)}(z) - e^{-|z|^2} \right| dA(z) \rightarrow 0 \quad \text{as } m \rightarrow \infty \quad \text{and } n \geq m\tau - M,$$

with uniform convergence for $z_0 \in K$. Equivalently, $\widehat{B}_{m,n}^{(z_0)} \rightarrow P$ as $m \rightarrow \infty$ and $n \geq m\tau - M$.

Proof. See Sect. 7. □

2.4. The expansion formula. Most of our results in this paper rely on a suitable approximation for $K_{m,n}(z, w)$ when z, w are both close to some point $z_0 \in X$, and m and n are large. We will here assume that Q be *real-analytic* in a neighbourhood of S_τ .

An adequate and rather far-reaching approximation formula was recently stated by Berman [3], p. 9, depending on methods from [6]; we will here only require a special case of his result. We will need to introduce some definitions.

For subsets $S \subset \mathbb{C}$ we write

$$\overline{\text{diag}}(S) = \{(z, \bar{z}) \in \mathbb{C}^2; z \in S\}.$$

That Q is real-analytic in a neighbourhood of S means that there exists a unique holomorphic function ψ defined in a neighbourhood of $\overline{\text{diag}}(S)$ in $\mathbb{C}^2$ such that

$$\psi(z, \bar{z}) = Q(z) \quad \text{for all } z \in \mathbb{C}.$$

Explicitly, one has

$$(2.6) \quad \psi(z + h, \overline{z + k}) = \sum_{i,j=0}^{\infty} [\partial^i \bar{\partial}^j Q](z) \frac{h^i \bar{k}^j}{i!j!}$$

when h and k are sufficiently close to zero. In particular, $[\partial^i \bar{\partial}^j \psi](z, \bar{z}) = \partial^i \bar{\partial}^j Q(z)$ for all $z \in \mathbb{C}$. Here, ∂_1 and ∂_2 denote differentiation w.r.t. the first and second coordinates.

Definition 2.7. Let b_0 and b_1 denote the functions

$$b_0 = \partial_1 \partial_2 \psi, \quad b_1 = \frac{1}{2} \partial_1 \partial_2 \log [\partial_1 \partial_2 \psi] = \frac{\partial_1^2 \partial_2^2 \psi \partial_1 \partial_2 \psi - \partial_1^2 \partial_2 \psi \partial_1 \partial_2^2 \psi}{2[\partial_1 \partial_2 \psi]^2}.$$

The function b_1 is well-defined where there is no division by 0, in particular in a neighborhood of $\text{diag}(X)$. Along the anti-diagonal, we have

$$b_0(z, \bar{z}) = \Delta Q(z) \quad \text{and} \quad b_1(z, \bar{z}) = \frac{1}{2} \Delta \log \Delta Q(z), \quad z \in X.$$

We note for later use that b_0 and b_1 are connected via

$$(2.7) \quad b_1(z, w) = \frac{1}{2} \frac{\partial}{\partial w} \left(\frac{1}{b_0(z, w)} \frac{\partial}{\partial z} b_0(z, w) \right).$$

We define the *first order approximating Bergman kernel* $K_m^1(z, w)$ by

$$K_m^1(z, w) = (mb_0(z, \bar{w}) + b_1(z, \bar{w})) e^{m\psi(z, \bar{w})}.$$

We have the following theorem.

Theorem 2.8. Assume that Q is real-analytic in $\mathbb{C}$. Let K be a compact subset of $S_\tau^\circ \cap X$. Fix a point $z_0 \in K$ and a number $M \geq 0$. Then there exists a number m_0 depending only on M and τ and a positive number ε depending only on K and M such that

$$|K_{m,n}(z, w) - K_m^1(z, w)| e^{-m(Q(z)+Q(w))/2} \leq Cm^{-1}, \quad z_0 \in K, \quad z, w \in \mathbb{D}(z_0; \varepsilon),$$

for all $m \geq m_0$ and $n \geq m\tau - M$. Here C is an absolute constant (depending only on Q). In particular, by restricting to the diagonal, we get

$$(2.8) \quad \left| K_{m,n}(z, z) e^{-mQ(z)} - \left(m\Delta Q(z) + \frac{1}{2} \Delta \log \Delta Q(z) \right) \right| \leq \frac{C}{m}, \quad z \in K,$$

when $m \geq m_0$ and $n \geq m\tau - M$.

Proof. See Sect. 6. See also Remark 2.9 below. □

Remark 2.9. We want to stress that Th. 2.8 has a long history; analogous expansions are well-known for weighted Bergman kernels, see, for instance, [13], [9], [19], [12], [6], [3], and the references therein. Moreover, as we mentioned above, Th. 2.8 is a slight modification of a more general result in several complex variables stated by Berman [3], Th. 3.8, which may be obtained by adapting the methods from [6]. In our proof, we make frequent use of ideas and techniques developed in [3], [5], [6], and in the book [22].

Remark 2.10. (Simple properties of K_m^1 .)

(1) Since b_0, b_1 and ψ are real on the anti-diagonal, a suitable version of the reflection principle implies that K_m^1 is Hermitian, that is, $\overline{K_m^1(z, w)} = K_m^1(w, z)$.

(2) Using the Taylor series for ψ at $(z, \bar{z})$ (see (2.6)) and ditto for Q and z , we get

$$(2.9) \quad 2 \operatorname{Re} \psi(z, \bar{w}) - Q(z) - Q(w) = -\Delta Q(w) |w - z|^2 + R(z, w),$$

where $R(z, w) = O(|w - z|^3)$ for z, w in a sufficiently small neighbourhood of z_0 , and the O is uniform for z_0 in a fixed compact subset of X . In case $\Delta Q(z_0) > 0$, it follows that

$$(2.10) \quad 2 \operatorname{Re} \psi(z, \bar{w}) - Q(z) - Q(w) \leq -\delta_0 |w - z|^2, \quad z, w \in \mathbb{D}(z_0; 2\varepsilon),$$

with $\delta_0 = \frac{1}{2} \Delta Q(z_0) > 0$, provided that the positive number ε is chosen small enough. More generally, if we fix a compact subset $K \Subset X$, and put

$$\delta_0 = \frac{1}{2} \inf_{\zeta \in K} \{\Delta Q(\zeta)\},$$

we may find an $\varepsilon > 0$ such that (2.10) holds for all $z_0 \in K$. With a perhaps somewhat smaller $\varepsilon > 0$, we may also ensure that the functions b_0 and b_1 are bounded and holomorphic in the set $\{(z, w); z, \bar{w} \in \mathbb{D}(z_0; 2\varepsilon), z_0 \in K\}$. We infer that

$$(2.11) \quad |K_m^1(z, w)|^2 e^{-m(Q(z)+Q(w))} \leq C m^2 e^{-m\delta_0 |z-w|^2}, \quad z, w \in \mathbb{D}(z_0; 2\varepsilon), \quad z_0 \in K,$$

with a number C depending only on K and ε .

2.5. Extensions and possible generalizations. We discuss possible extensions of Th. 2.8 (and also of Th. 2.6). Again, these extensions (at least of Th. 2.8) are essentially implied by a result [[3], Th. 3.8], stated by R. Berman.

For a given positive integer k , one may define a k -th order approximating Bergman kernel

$$K_m^k(z, w) = (mb_0(z, \bar{w}) + b_1(z, \bar{w}) + \dots + m^{-k+1}b_k(z, \bar{w})) e^{m\psi(z, \bar{w})},$$

for z and w in a neighbourhood of $\overline{\operatorname{diag}}(X)$ where b_i are certain holomorphic functions defined in a neighbourhood of $\overline{\operatorname{diag}}(X)$. It can be shown that for $z_0 \in X$ there exists $\varepsilon > 0$ such that

$$|K_{m,n}(z, w) - K_m^k(z, w)|^2 e^{-m(Q(z)+Q(w))} \leq C m^{-2k},$$

whenever $z, w \in \mathbb{D}(z_0; \varepsilon)$ and $n \geq m\tau - 1$. The coefficients b_i can in principle be determined from a recursion formula involving partial differential equations of increasing order, compare with [Berman et al. [6], (2.15), p.9], where a closely related formula is given. However, the analysis required for calculating higher order coefficients b_i for $i \geq 2$ seems to be quite involved, and the first order approximation seems to be sufficient for many practical purposes, cf. [1]. We therefore prefer a more direct approach here.

Th. 2.8 can also be generalized in another direction – one may relax the assumption that Q be real-analytic, and instead assume that e.g. $Q \in C^\infty(\mathbb{C})$. In this case, the functions b_i and ψ will be almost-holomorphic at the anti-diagonal (cf. e.g. [9] for a relevant discussion of such functions). The modifications needed for proving Th. 2.8 in this more general case are based on standard arguments; they are essentially as outlined in [[6], Subsect. 2.6 and p. 15]. We leave the details to the interested reader. However, the reader should note that, using this generalized version of Th. 2.8, one may easily extend our proof of Th. 2.6, to the case of C^∞ -smooth weights.

2.6. The Bargmann–Fock case and harmonic measure. When $z_0 \in \mathbb{C} \setminus (\mathcal{S}_\tau^\circ \cap X)$, Prop. 2.2 yields that the measures $B_{m,n}^{(z_0)}$ tend to concentrate on $\mathcal{S}_\tau$ as $m \rightarrow \infty$ and $n - m\tau \rightarrow 0$. However our general results provide no further information regarding the asymptotic distribution.

We now specialize to the Bargmann–Fock weight $Q(z) = |z|^2$. We then obviously have $X = \mathbb{C}$. It will be convenient to introduce the functions (truncated exponentials)

$$E_k(z) = \sum_{j=0}^k \frac{z^j}{j!}.$$

It is then easy to check that $K_{m,n}(z, w) = mE_{n-1}(mz\bar{w})$, $\mathcal{S}_\tau = \overline{\mathbb{D}}(0; \sqrt{\tau})$, and $\widehat{Q}_\tau(z) = \tau + \tau \log(|z|^2/\tau)$ when $|z|^2 \geq \tau$. We infer that

$$(2.12) \quad dB_{m,n}^{(z_0)}(z) = m \frac{|E_{n-1}(mz\bar{z}_0)|^2}{E_{n-1}(m|z_0|^2)} e^{-m|z|^2} dA(z).$$

Let $\mathbb{C}^* = \mathbb{C} \cup \{\infty\}$ denote the extended plane.

Theorem 2.11. *Let $Q(z) = |z|^2$ and $z_0 \in \mathbb{C} \setminus \mathcal{S}_\tau$. Then, as $m \rightarrow \infty$ and $n/m \rightarrow \tau$, the measures $B_{m,n}^{(z_0)}$ converge to harmonic measure at z_0 with respect to $\mathbb{C}^* \setminus \mathcal{S}_\tau$.*

Proof. See Sect. 9. □

Remark 2.12. The above theorem is a special case of Theorem 8.7 in [1].

2.7. Weighted L^2 estimates for the $\bar{\partial}$ -equation with a growth constraint. Our analysis depends in an essential way on having a good control for the norm-minimal solution u_* in the space $L_{mQ,n}^2$ to the $\bar{\partial}$ -equation $\bar{\partial}u = f$, where f is a given suitable test-function such that $\text{supp } f \subset \mathcal{S}_\tau \cap X$. We would not have a problem if u_* were just required to be of class L_{mQ}^2 (and not $L_{mQ,n}^2$), for then an elementary version of the well-known L^2 estimates of Hörmander apply, see e.g. [18] or [17]. The requirement that u_* be of restricted growth gives rise to some difficulties, but we shall see in Sect. 4 (notably Th. 4.1) that the classical estimates can be adapted to our situation. (Estimates very similar to ours were given independently by Berman in [4] after this note was completed.)

3. PREPARATORY LEMMAS

3.1. Preliminaries. In this section, we prove Prop. 2.2. At the same time we will get the opportunity to display some elementary inequalities which will be used in later sections. For the results obtained here, it is enough to suppose that Q is $C^{1,1}$ -smooth (and satisfies the growth assumption (1.5)). Thus ΔQ makes sense almost everywhere and is of class $L_{\text{loc}}^\infty(\mathbb{C})$.

3.2. Subharmonicity estimates. We start by giving two simple lemmas which are based on the sub-mean property of subharmonic functions.

Lemma 3.1. *Let ϕ be a weight of class $C^{1,1}(\mathbb{C})$ and put $A = \text{ess sup } \{\Delta\phi(\zeta); \zeta \in \mathbb{D}\}$. Then if u is bounded and holomorphic in $\mathbb{D}$, we have that*

$$|u(0)|^2 e^{-\phi(0)} \leq \int_{\mathbb{D}} |u(\zeta)|^2 e^{-\phi(\zeta)} e^{A|\zeta|^2} dA(\zeta) \leq e^A \int_{\mathbb{D}} |u(\zeta)|^2 e^{-\phi(\zeta)} dA(\zeta).$$

Proof. Consider the function $F(\zeta) = |u(\zeta)|^2 e^{-\phi(\zeta) + A|\zeta|^2}$, which satisfies

$$\Delta \log F = \Delta \log |u|^2 - \Delta\phi + A \geq 0 \quad \text{a.e. on } \mathbb{D},$$

making F logarithmically subharmonic. But then F is subharmonic itself, and so

$$F(0) \leq \int_{\mathbb{D}} F(z) dA(z).$$

The assertion of the lemma is immediate from this. □

Lemma 3.2. Let $Q \in C^{1,1}(\mathbb{C})$ and fix $\delta > 0$ and $z \in \mathbb{C}$ and put $A = A_{z,\delta} = \text{ess sup } \{\Delta Q(\zeta); \zeta \in \mathbb{D}(z; \delta)\}$. Also, let u be holomorphic and bounded in $\mathbb{D}(z; \delta/\sqrt{m})$. Then

$$|u(z)|^2 e^{-mQ(z)} \leq m\delta^{-2} e^{A\delta^2} \int_{\mathbb{D}(z; \delta/\sqrt{m})} |u(\zeta)|^2 e^{-mQ(\zeta)} dA(\zeta).$$

Proof. The assertion follows if we make the change of variables $\zeta = z + \delta\xi/\sqrt{m}$ where $\zeta \in \mathbb{D}(z; \delta/\sqrt{m})$ and $\xi \in \mathbb{D}$, and apply Lemma 3.1 with the weight $\phi(\xi) = mQ(\zeta)$. $\square$

We note the following consequence of the subharmonicity estimates. We will frequently need it in later sections.

Lemma 3.3. Let K be a compact subset of $\mathbb{C}$ and δ a given positive number. We put

$$K_\delta = \{z \in \mathbb{C}; \text{dist}(z, K) \leq \delta\} \quad \text{and} \quad A = \text{ess sup } \{\Delta Q(z); z \in K_\delta\}.$$

Then, for all $m, n \geq 1$,

$$|K_{m,n}(z, w)|^2 e^{-mQ(z)} \leq m\delta^{-2} e^{\delta^2 A} \int_{\mathbb{D}(z; \delta/\sqrt{m})} |K_{m,n}(\zeta, w)|^2 e^{-mQ(\zeta)} dA(\zeta), \quad z \in K, \quad w \in \mathbb{C}.$$

Proof. Apply Lemma 3.2 to $u(\zeta) = K_{m,n}(\zeta, w)$. $\square$

3.3. A weak maximum principle for weighted polynomials. Maximum principles for weighted polynomials have a long history, see e.g. [21], Chap. III. The following simple lemma will suffice for our present purposes; it is a consequence of [21], Th. III.2.1. (Recall that $\mathcal{S}_\tau$ denotes the coincidence set $\{Q = \widehat{Q}_\tau\}$, see (2.2).)

Lemma 3.4. Suppose that a polynomial u of degree at most $n - 1$ satisfies $|u(z)|^2 e^{-mQ(z)} \leq 1$ on $\mathcal{S}_{\tau(m,n)}$, where $\tau(m, n) = (n - 1)/m < \rho$. Then $|u(z)|^2 e^{-m\widehat{Q}_{\tau(m,n)}(z)} \leq 1$ on $\mathbb{C}$.

Proof. Put $q(z) = m^{-1} \log |u(z)|^2$. The assumptions on u imply that $q \in \text{SH}_{\tau(m,n)}$ and that $q \leq Q$ on $\mathcal{S}_{\tau(m,n)}$. Hence $q \leq \widehat{Q}_{\tau(m,n)}$ on $\mathbb{C}$, as desired. $\square$

Lemma 3.5. Let $u \in H_{m,n}$, and suppose that $n \leq m\tau + 1$. Then

$$|u(z)|^2 \leq m e^A \|u\|_{mQ}^2 e^{m\widehat{Q}_\tau(z)},$$

where A denotes the essential supremum of ΔQ over the set $\{z \in \mathbb{C}; \text{dist}(z, \mathcal{S}_\tau) \leq 1\}$.

Proof. The assertion that $n \leq m\tau + 1$ is equivalent to that $\tau(m, n) \leq \tau$ where $\tau(m, n) = (n - 1)/m$. Thus $\mathcal{S}_{\tau(m,n)} \subset \mathcal{S}_\tau$ and $\widehat{Q}_{\tau(m,n)} \leq \widehat{Q}_\tau$. An application of Lemma 3.2 with $\delta = 1$ now gives

$$|u(z)|^2 e^{-mQ(z)} \leq m e^A \int_{\mathbb{D}(z; 1)} |u(\zeta)|^2 e^{-mQ(\zeta)} dA(\zeta) \leq m e^A \int_{\mathbb{C}} |u(\zeta)|^2 e^{-mQ(\zeta)} dA(\zeta), \quad z \in \mathcal{S}_\tau.$$

As a consequence, the same estimate holds on $\mathcal{S}_{\tau(m,n)}$. We can thus apply Lemma 3.4. It yields that

$$|u(z)|^2 \leq m e^A \|u\|_{mQ}^2 e^{m\widehat{Q}_{\tau(m,n)}(z)}, \quad z \in \mathbb{C}.$$

The desired assertion follows, since $\widehat{Q}_{\tau(m,n)} \leq \widehat{Q}_\tau$. $\square$

3.4. The proof of Proposition 2.2. Fix two points z and z_0 in $\mathbb{C}$. We apply Lemma 3.5 to the polynomial

$$u(\zeta) = \frac{K_{m,n}(\zeta, z_0)}{\sqrt{K_{m,n}(z_0, z_0)}},$$

which is of class $H_{m,n}$ and satisfies $\|u\|_{mQ} = 1$. It yields that $|u(z)|^2 \leq m e^A e^{m\widehat{Q}_\tau(z)}$, or

$$(3.1) \quad \mathcal{B}_{m,n}^{(w)}(z) = |u(z)|^2 e^{-mQ(z)} \leq m e^A e^{m(\widehat{Q}_\tau(z) - Q(z))}, \quad z \in \mathbb{C}, \quad n \leq m\tau + 1.$$

Now let Λ be an open neighborhood of $\mathcal{S}_\tau$. Since $Q > \widehat{Q}_\tau$ on $\mathbb{C} \setminus \mathcal{S}_\tau$, the continuity of the functions involved coupled with the growth conditions (2.3) and (1.5) yield that $Q - \widehat{Q}_\tau$ is bounded below by a positive number on $\mathbb{C} \setminus \Lambda$. It follows that

$$B_{m,n}^{(z_0)}(\mathbb{C} \setminus \Lambda) = \int_{\mathbb{C} \setminus \Lambda} B_{m,n}^{(z_0)} dA \rightarrow 0$$

as $m \rightarrow \infty$ and $n \leq m\tau + 1$. Since $B_{m,n}^{(z_0)}$ is a p.m. on $\mathbb{C}$, the assertion of Prop. 2.2 follows. $\square$

3.5. An estimate for the one-point function. Our proof of Prop. 2.2 implies a useful estimate for the one-point function $z \mapsto K_{m,n}(z, z)e^{-mQ(z)}$. The following result is implicit in [3].

Proposition 3.6. *Suppose that $n \leq m\tau + 1$. Then there exists a number C depending only on τ such that*

$$(3.2) \quad K_{m,n}(z, z)e^{-mQ(z)} \leq Cme^{-m(Q(z) - \widehat{Q}_\tau(z))}, \quad z \in \mathbb{C},$$

$$(3.3) \quad |K_{m,n}(z, w)|^2 e^{-m(Q(z) + Q(w))} \leq Cm^2 e^{-m(Q(z) - \widehat{Q}_\tau(z))} e^{-m(Q(w) - \widehat{Q}_\tau(w))}, \quad z, w \in \mathbb{C}.$$

In particular, $|K_{m,n}(z, w)|^2 e^{-m(Q(z) + Q(w))} \leq Cm^2$ for all z and w .

Proof. The function $B_{m,n}^{(z_0)}(z)$ in the diagonal case $z = z_0$ reduces to

$$B_{m,n}^{(z)}(z) = K_{m,n}(z, z)e^{-mQ(z)}.$$

Thus the estimate (3.1) implies

$$K_{m,n}(z, z)e^{-mQ(z)} \leq me^A e^{m(\widehat{Q}_\tau(z) - Q(z))},$$

which proves (3.2). In order to get (3.3) it now suffices to use the Cauchy–Schwarz inequality $|K_{m,n}(z, w)|^2 \leq K_{m,n}(z, z)K_{m,n}(w, w)$ and apply (3.2) to the two factors in the right hand side. $\square$

3.6. Cut-off functions. We will in the following frequently use cut-off functions with various properties. For convenience of the reader who may not be familiar with these details, we collect the necessary facts here.

Given any numbers $\delta > 0$, $r > 0$ and $C > 1$, there exists $\chi \in C_0^\infty(\mathbb{C})$ such that $\chi = 1$ on $\mathbb{D}(0; \delta)$, $\chi = 0$ outside $\mathbb{D}(0; \delta(1 + r))$, $\chi \leq 1$ and $|\bar{\partial}\chi|^2 \leq (C/r^2)\chi$ on $\mathbb{C}$.² Moreover, with such a choice for χ , it follows that $|\bar{\partial}\chi|^2 \leq (C/r^2)\delta^{-2}\chi$ on $\mathbb{C}$. It is then easy to check that $\|\bar{\partial}\chi\|_{L^2}^2 \leq C(1 + 2/r)$.

Later on, we will often use the values $\delta = 3\varepsilon/2$ and $\delta(1 + r) = 2\varepsilon$, where $\varepsilon > 0$ is given. We may then arrange that $\|\bar{\partial}\chi\|_{L^2} \leq 3$.

4. WEIGHTED ESTIMATES FOR THE $\bar{\partial}$ -EQUATION WITH A GROWTH CONSTRAINT

4.1. General introduction; the Bergman projection and the $\bar{\partial}$ -equation. Let ϕ be a weight on $\mathbb{C}$ (cf. Subsect. 1.3). We assume throughout that ϕ is of class $C^{1,1}(\mathbb{C})$ (so that $\Delta\phi \in L_{\text{loc}}^\infty(\mathbb{C})$), and that $\int_{\mathbb{C}} e^{-\phi} dA < \infty$ (so that L_ϕ^2 contains the constant functions). Also fix a positive integer n and recall the definition of the "truncated" spaces $L_{\phi,n}^2$ and $A_{\phi,n}^2$ (see (1.3) and (1.4)).

Let $K_{\phi,n}$ denote the reproducing kernel for $A_{\phi,n}^2$, and let $P_{\phi,n} : L_\phi^2 \rightarrow A_{\phi,n}^2$ be the orthogonal projection,

$$(4.1) \quad P_{\phi,n}u(w) = \int_{\mathbb{C}} u(z)K_{\phi,n}(w, z)e^{-\phi(z)} dA(z), \quad u \in L_\phi^2.$$

We will in later sections frequently need to estimate $P_{\phi,n}u(w)$, especially when u is holomorphic in a neighbourhood of w .

²Such a χ can be obtained by standard regularization applied to the Lipschitzian function which equals $(1 - (\delta^{-1}|z| - 1)/r)^2$ for $1 \leq |z| \leq r$, and is otherwise locally constant on $\mathbb{C}$.

Now note that for f in the class $C_0(\mathbb{C})$ (continuous functions with compact support), the Cauchy transform $u = Cf$ given by

$$Cf(w) = \int_{\mathbb{C}} \frac{f(z)}{w - z} dA(z),$$

satisfies the $\bar{\partial}$ -equation

$$(4.2) \quad \bar{\partial}u = f.$$

Moreover, $u = Cf$ is bounded, and is therefore of class $L^2_{\phi,n}$ for any $n \geq 1$. Thus the function

$$(4.3) \quad u_*(w) = u(w) - P_{\phi,n}u(w),$$

solves (4.2) and is of class $L^2_{\phi,n}$. It is easy to verify that u_* defined by (4.3) is the unique norm-minimal solution to (4.2) in $L^2_{\phi,n}$ whenever $u \in L^2_{\phi,n}$ is a solution to (4.2). Hence the study of the orthogonal projection $P_{\phi,n}u$ is equivalent to the study of the $L^2_{\phi,n}$ -minimal solution u_* to (4.2).

It is useful to observe that u_* is characterized amongst the solutions of class $L^2_{\phi,n}$ to (4.2) by the condition $u_* \perp A^2_{\phi,n}$, or

$$(4.4) \quad \int_{\mathbb{C}} u_*(z) \overline{h(z)} e^{-\phi(z)} dA(z) = 0, \quad \text{for all } h \in A^2_{\phi,n}.$$

Our principal result in this section, Th. 4.1, states that a variant of the elementary one-dimensional form of the L^2 estimates of Hörmander are valid for u_* . The important results for the further developments in this paper are however Th. 4.4 and Cor. 4.5. The reader may consider to glance at those results and skip to the next section, at a first read.

4.2. L^2 estimates. The L^2 estimates of Hörmander in the most elementary, one-dimensional form applies only to weights ϕ which are strictly subharmonic in the entire plane $\mathbb{C}$. The result states that u_0 , the L^2_{ϕ} -minimal solution to (4.2) (where $f \in C_0(\mathbb{C})$) satisfies

$$(4.5) \quad \int_{\mathbb{C}} |u_0|^2 e^{-\phi} dA \leq \int_{\mathbb{C}} |f|^2 \frac{e^{-\phi}}{\Delta\phi} dA,$$

provided that ϕ is C^2 -smooth on $\mathbb{C}$. See [18], eq. (4.2.6), p. 250 (this is essentially just Green's formula).

It is important to observe that the estimate (4.5) remains valid for strictly subharmonic weights ϕ in the larger class $C^{1,1}(\mathbb{C})$ (that ϕ is strictly subharmonic then means that $\Delta\phi > 0$ a.e. on $\mathbb{C}$). The proof in [18], Subsect. 4.2 goes through without changes in this more general case.

4.3. Weighted $L^2_{\phi,n}$ estimates. We have the following theorem, where we consider two weights ϕ and $\widehat{\phi}$ with various properties. The practically minded reader may, with little loss of generality, think of $\phi = mQ$, $\widehat{\phi}(z) = m\widehat{Q}_{\tau}(z) + \varepsilon \log(1 + |z|^2)$, and $\Sigma = S_{\tau}$. This will essentially be the case in all our later applications.

Theorem 4.1. *Let Σ be a compact subset of $\mathbb{C}$, ϕ , $\widehat{\phi}$ and ϱ three real-valued functions of class $C^{1,1}(\mathbb{C})$, and n a positive integer. We assume that:*

- (1) $L^2_{\widehat{\phi}}$ contains the constants and there are non-negative numbers α and β such that

$$(4.6) \quad \widehat{\phi} \leq \phi + \alpha \quad \text{on } \mathbb{C} \quad \text{and} \quad \phi \leq \beta + \widehat{\phi} \quad \text{on } \Sigma,$$

- (2) $A^2_{\widehat{\phi}} \subset A^2_{\phi,n}$,

- (3) The function $\widehat{\phi} + \varrho$ is strictly subharmonic on $\mathbb{C}$,

- (4) ϱ is locally constant on $\mathbb{C} \setminus \Sigma$,

(5) There exists a number κ , $0 < \kappa < 1$, such that

$$(4.7) \quad \frac{|\bar{\partial}\varrho|^2}{\widehat{\Delta\phi} + \Delta\varrho} \leq \frac{\kappa^2}{e^{\alpha+\beta}} \quad \text{a.e. on } \Sigma.$$

Let $f \in C_0^\infty(\mathbb{C})$ be such that

$$\text{supp } f \subset \Sigma.$$

Then u_* , the $L_{\phi,n}^2$ -minimal solution to $\bar{\partial}u = f$ satisfies

$$(4.8) \quad \int_{\mathbb{C}} |u_*|^2 e^{\varrho-\phi} dA \leq \frac{e^{\alpha+\beta}}{(1-\kappa)^2} \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho-\phi}}{\widehat{\Delta\phi} + \Delta\varrho} dA.$$

Before we give the proof, we note a simple lemma, which will be put to repeated use.

Lemma 4.2. Suppose that $f \in C_0(\mathbb{C})$. Then (4.2) has a solution u in L_{ϕ}^2 . Moreover v_* , the L_{ϕ}^2 -minimal solution to (4.2) is of class $L_{\phi,n}^2$.

Proof. The assumptions imply that the Cauchy transform Cf is in L_{ϕ}^2 . Thus the L_{ϕ}^2 -minimal solution v_* to (4.2) exists, necessarily of the form $v_* = Cf + g$ with some $g \in A_{\phi}^2$. In view of (3), we know that $g \in A_{\phi,n}^2$. The assertion is now immediate. $\square$

Proof of Theorem 4.1. Assume that the right hand side in (4.8) is finite. In view of (4.4), the condition that u_* is $L_{\phi,n}^2$ -minimal may be expressed as

$$\int_{\mathbb{C}} u_* e^{\varrho} \bar{h} e^{-(\phi+\varrho)} dA = 0 \quad \text{for all } h \in A_{\phi,n}^2.$$

The latter relation means that the function $w_0 = u_* e^{\varrho}$ minimizes the norm in $L_{\phi+\varrho}^2$ over elements of the (non-closed) subspace

$$e^{\varrho} \cdot L_{\phi,n}^2 = \{w; \quad w = h e^{\varrho}, \quad \text{where } h \in L_{\phi,n}^2\} \subset L_{\phi+\varrho}^2$$

which solve the (modified) $\bar{\partial}$ -equation

$$(4.9) \quad \bar{\partial}w = \bar{\partial}(u_* e^{\varrho}) = f e^{\varrho} + u_* \bar{\partial}(e^{\varrho}).$$

Now, since ϱ is bounded on $\mathbb{C}$ and locally constant outside Σ , we have

$$L_{\phi+\varrho,n}^2 = e^{\varrho} \cdot L_{\phi,n}^2 = L_{\phi,n}^2 \quad (\text{as sets}),$$

and we conclude that w_0 is the norm-minimal solution to (4.9) in $L_{\phi+\varrho,n}^2$.

Let v_* denote the L_{ϕ}^2 -minimal solution to $\bar{\partial}u = f$ (see Lemma 4.2). We form the continuous function $g = \bar{\partial}((u_* - v_*)e^{\varrho}) = (u_* - v_*)\bar{\partial}(e^{\varrho})$, whose support is contained in Σ , and consider the $\bar{\partial}$ -equation

$$(4.10) \quad \bar{\partial}\xi = g = (u_* - v_*)\bar{\partial}(e^{\varrho}).$$

The assertion of Lemma 4.2 remains valid if $\widehat{\phi}$ is replaced by $\widehat{\phi} + \varrho$; it follows that (4.10) has a solution $\xi \in L_{\phi+\varrho}^2$, and moreover, if ξ_0 denotes the norm-minimal solution to (4.10) in $L_{\phi+\varrho}^2$, we have $\xi_0 \in L_{\phi+\varrho,n}^2$.

We now continue by forming the function

$$w_1 = v_* e^{\varrho} + \xi_0.$$

It is clear that $w_1 \in L_{\phi+\varrho,n}^2$, and a calculation yields that

$$(4.11) \quad \bar{\partial}w_1 = f e^{\varrho} + v_* \bar{\partial}(e^{\varrho}) + (u_* - v_*)\bar{\partial}(e^{\varrho}) = \bar{\partial}(u_* e^{\varrho}) = \bar{\partial}w_0.$$

Since w_0 is norm-minimal in $L^2_{\widehat{\phi}+\varrho,n}$ over functions w such that $\bar{\partial}w = \bar{\partial}w_0$, we must have

$$(4.12) \quad \int_{\mathbb{C}} |w_0|^2 e^{-(\widehat{\phi}+\varrho)} dA \leq \int_{\mathbb{C}} |w_1|^2 e^{-(\widehat{\phi}+\varrho)} dA \leq e^\alpha \int_{\mathbb{C}} |w_1|^2 e^{-(\widehat{\phi}+\varrho)} dA,$$

where we have used the condition $\widehat{\phi} \leq \phi + \alpha$ to deduce the second inequality. Moreover, since ξ_0 is norm-minimal in $L^2_{\widehat{\phi}+\varrho}$ amongst solutions to (4.10), we have

$$\int_{\mathbb{C}} w_1 \bar{h} e^{-(\widehat{\phi}+\varrho)} dA = \int_{\mathbb{C}} v_* \bar{h} e^{-\widehat{\phi}} dA + \int_{\mathbb{C}} \xi_0 \bar{h} e^{-(\widehat{\phi}+\varrho)} dA = 0$$

for all $h \in A^2_{\widehat{\phi}+\varrho}$, so the function w_1 is in fact the norm-minimal solution to a $\bar{\partial}$ -equation in $L^2_{\widehat{\phi}+\varrho}$.

The $\bar{\partial}$ -equation satisfied by w_1 is (see (4.11))

$$\bar{\partial}w_1 = f e^\varrho + u_* \bar{\partial}(e^\varrho) = f e^\varrho + u_* e^\varrho \bar{\partial}\varrho.$$

By the estimate (4.5) applied to the weight $\widehat{\phi} + \varrho$, we obtain that

$$(4.13) \quad \int_{\mathbb{C}} |w_1|^2 e^{-(\widehat{\phi}+\varrho)} dA \leq \int_{\mathbb{C}} |f e^\varrho + u_* e^\varrho \bar{\partial}\varrho|^2 \frac{e^{-(\widehat{\phi}+\varrho)}}{\Delta(\widehat{\phi} + \varrho)} dA = \int_{\mathbb{C}} |f + u_* \bar{\partial}\varrho|^2 \frac{e^{\varrho-\widehat{\phi}}}{\Delta\widehat{\phi} + \Delta\varrho} dA.$$

Since f and $\bar{\partial}\varrho$ are supported in Σ , and since $e^{-\widehat{\phi}} \leq e^\beta e^{-\phi}$ there (see (4.6)), it is seen that the right hand side in (4.13) can be estimated by

$$(4.14) \quad e^\beta \int_{\mathbb{C}} |f + u_* \bar{\partial}\varrho|^2 \frac{e^{\varrho-\phi}}{\Delta\widehat{\phi} + \Delta\varrho} dA.$$

For $t > 0$ we now use the inequality $|a + b|^2 \leq (1+t)|a|^2 + (1+t^{-1})|b|^2$ and the condition (4.7) to conclude that the integral in (4.14) is dominated by

$$(4.15) \quad \begin{aligned} (1+t) \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho-\phi}}{\Delta\widehat{\phi} + \Delta\varrho} dA + (1+t^{-1}) \int_{\mathbb{C}} |u_*|^2 \frac{|\bar{\partial}\varrho|^2}{\Delta\widehat{\phi} + \Delta\varrho} e^{\varrho-\phi} dA &\leq \\ &\leq (1+t) \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho-\phi}}{\Delta\widehat{\phi} + \Delta\varrho} dA + (1+t^{-1}) \frac{\kappa^2}{e^{\alpha+\beta}} \int_{\mathbb{C}} |u_*|^2 e^{\varrho-\phi} dA. \end{aligned}$$

Tracing back through (4.12), (4.13), (4.14) and (4.15), we get

$$\int_{\mathbb{C}} |u_*|^2 e^{\varrho-\phi} dA \leq e^{\alpha+\beta} (1+t) \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho-\phi}}{\Delta\widehat{\phi} + \Delta\varrho} dA + (1+t^{-1}) \kappa^2 \int_{\mathbb{C}} |u_*|^2 e^{\varrho-\phi} dA,$$

which we write as

$$(1 - (1+t^{-1})\kappa^2) \int_{\mathbb{C}} |u_*|^2 e^{\varrho-\phi} dA \leq e^{\alpha+\beta} (1+t) \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho-\phi}}{\Delta\widehat{\phi} + \Delta\varrho} dA.$$

The desired estimate (4.8) now follows if we pick $t = \kappa/(1 - \kappa)$. $\square$

4.4. Implementation scheme. We now fix $Q \in C^{1,1}(\mathbb{C})$ satisfying the growth assumption (1.5) with a fixed $\rho > 0$. Adding a constant to Q does not change the problem and so we may assume that $Q \geq 1$ on $\mathbb{C}$. Let us put

$$(4.16) \quad q_\tau = \sup_{z \in \mathcal{S}_\tau} \{Q(z)\}.$$

We next fix a positive number $\tau < \rho$ and two positive numbers M_0 and M_1 such that

$$(4.17) \quad M_1 \log(1 + |z|^2) \leq M_0 \widehat{Q}_\tau(z), \quad z \in \mathbb{C}.$$

This is possible since $\widehat{Q}_\tau \geq 1$ and $\widehat{Q}_\tau(z) = \tau \log |z|^2 + O(1)$ when $z \rightarrow \infty$ (see (2.3)). In particular, it yields that $M_1 \leq M_0 \tau$. We now define

$$(4.18) \quad \phi_m = mQ \quad \text{and} \quad \widehat{\phi}_{m,M_0,M_1}(z) = (m - M_0)\widehat{Q}_\tau(z) + M_1 \log(1 + |z|^2).$$

Note that $\widehat{\phi}_{m,M_0,M_1}$ is strictly subharmonic on $\mathbb{C}$ whenever $m \geq M_0$ with

$$(4.19) \quad \Delta \widehat{\phi}_{m,M_0,M_1}(z) = (m - M_0)\Delta \widehat{Q}_\tau(z) + M_1 (1 + |z|^2)^{-2} \geq M_1 (1 + |z|^2)^{-2},$$

and that (4.17) implies (since $\widehat{Q}_\tau \leq Q$)

$$\widehat{\phi}_{m,M_0,M_1} \leq \phi_m \quad \text{on } \mathbb{C}.$$

Furthermore, (2.3) implies that

$$(4.20) \quad \widehat{\phi}_{m,M_0,M_1} = ((m - M_0)\tau + M_1) \log |z|^2 + O(1), \quad \text{as } z \rightarrow \infty.$$

It yields that

$$\phi_m(z) - \widehat{\phi}_{m,M_0,M_1}(z) \geq m(\rho - \tau) \log |z|^2 + O(1), \quad \text{as } z \rightarrow \infty.$$

Note also that

$$A_{\phi_m,n}^2 = A_{mQ}^2 \cap \mathcal{P}_n = H_{m,n}.$$

We now check the conditions (1) and (2) of Th. 4.1. (Recall that $\lfloor x \rfloor$ denotes the largest integer which is strictly smaller than x .)

Lemma 4.3. *Condition (1) holds for $\phi = \phi_m$, $\widehat{\phi} = \widehat{\phi}_{m,M_0,M_1}$, $\Sigma = \mathcal{S}_\tau$, $\alpha = 0$ and $\beta = M_0 q_\tau$, provided that $(m - M_0)\tau + M_1 > 1$. Condition (2), i.e. $A_{\phi_m,M_0,M_1}^2 \subset H_{m,n}$, holds if $n \geq \lfloor (m - M_0)\tau + M_1 \rfloor$. Thus conditions (1) and (2) hold whenever*

$$(4.21) \quad n \geq \lfloor (m - M_0)\tau + M_1 \rfloor > 0.$$

Proof. We have already shown that $\widehat{\phi}_{m,M_0,M_1} \leq \phi_m$ on $\mathbb{C}$. Moreover $\phi_m \leq \beta + \widehat{\phi}_{m,M_0,M_1}$ on $\mathcal{S}_\tau$, since $Q = \widehat{Q}_\tau$ there. The assertion that $\widehat{\phi}_{m,M_0,M_1} \leq \phi_m$ implies that $A_{\phi_m,M_0,M_1}^2 \subset A_{\phi_m}^2$, and it remains only to prove that $A_{\phi_m,M_0,M_1}^2 \subset \mathcal{P}_n$. But this follows from (4.20), (4.21), and the fact that $\int_{\mathbb{C}} (1 + |z|^2)^{-r} dA(z) < \infty$ if and only if $r > 1$. $\square$

We now apply Th. 4.1. It will be convenient to define

$$(4.22) \quad c_\tau = \inf_{z \in \mathcal{S}_\tau} \left\{ (1 + |z|^2)^{-2} \right\}.$$

Theorem 4.4. *Let $Q \in C^{1,1}(\mathbb{C})$ and $Q \geq 1$ on $\mathbb{C}$. Fix two positive numbers M_0 and M_1 such that relation (4.17) is satisfied, and define the positive numbers q_τ and c_τ by (4.16) and (4.22), and let $\widehat{\phi}_{m,M_0,M_1}$ be defined by (4.18). Suppose there are real-valued functions $\varrho_m \in C^{1,1}(\mathbb{C})$ which are locally constant on $\mathbb{C} \setminus \mathcal{S}_\tau$ such that $\widehat{\phi}_{m,M_0,M_1} + \varrho_m$ is strictly subharmonic on $\mathbb{C}$ and that for some positive number $\kappa < 1$ we have*

$$\frac{|\bar{\partial} \varrho_m|^2}{\Delta \widehat{\phi}_{m,M_0,M_1} + \Delta \varrho_m} \leq \frac{\kappa^2}{e^{M_0 q_\tau}} \quad \text{a.e. on } \mathbb{C}.$$

Suppose furthermore that $f \in C_0^\infty(\mathbb{C})$ satisfies

$$\text{supp } f \subset \mathcal{S}_\tau.$$

Then, if $n \geq \lfloor (m - M_0)\tau + M_1 \rfloor > 0$, we have that u_* , the $L_{mQ,n}^2$ -minimal solution to $\bar{\partial} u = f$, satisfies

$$\int_{\mathbb{C}} |u_*|^2 e^{\varrho_m - mQ} dA \leq \frac{e^{M_0 q_\tau}}{(1 - \kappa)^2} \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho_m - mQ}}{(m - M_0)\Delta Q + \Delta \varrho_m + M_1 c_\tau} dA.$$

Proof. All conditions in Th. 4.1 are fulfilled with $\phi = mQ$, $\widehat{\phi} = \widehat{\phi}_{m,M_0,M_1}$, $Q = Q_m$, $\alpha = 0$ and $\beta = M_0q_\tau$. In view of (4.19), it yields that

$$\begin{aligned} \int_{\mathbb{C}} |u_*|^2 e^{\varrho_m - mQ} dA &\leq \frac{e^{M_0q_\tau}}{(1-\kappa)^2} \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho_m - mQ}}{\Delta \widehat{\phi}_{m,M_0,M_1} + \Delta Q_m} dA \leq \\ &\leq \frac{e^{M_0q_\tau}}{(1-\kappa)^2} \int_{\mathbb{C}} |f|^2 \frac{e^{\varrho_m - mQ}}{(m - M_0)\Delta \widehat{Q}_\tau + \Delta Q_m + M_1c_\tau} dA. \end{aligned}$$

The assertion is now immediate from (4.4), since $\Delta Q = \Delta \widehat{Q}_\tau$ a.e. on $\text{supp } f$. $\square$

In Sect. 8, we shall use the full force of Th. 4.4. As for now, we just mention the following simple consequence, which also can be proved easily by more elementary means, cf. [3], p. 10.

Corollary 4.5. *Let $Q \in C^{1,1}(\mathbb{C})$ and $Q \geq 1$ on $\mathbb{C}$. Further, let Σ be a compact subset of $\mathcal{S}_\tau$. Put*

$$a = \text{ess inf } \{\Delta Q(z); z \in \Sigma\}.$$

Let $m_0 = \max\{2M_0, (1 + M_0)/\tau\}$ and assume that $f \in C_0^\infty(\mathbb{C})$ satisfies

$$\text{supp } f \subset \Sigma,$$

and that $n \geq (m - M_0)\tau + M_1$ and $m \geq m_0$. Then u_ , the $L^2_{mQ,n}$ -minimal solution to $\bar{\partial}u = f$ satisfies*

$$\|u_*\|_{mQ}^2 \leq \frac{2e^{M_0q_\tau}}{am + M_1c_\tau} \|f\|_{mQ}^2.$$

Proof. Take $Q_m = 0$ in Th. 4.4 and observe that $m - M_0 \geq m/2$ and also $(m - M_0)\tau + M_1 > 0$ whenever $m \geq m_0$. (Any other m_0 having these properties would work as well, of course.) $\square$

5. APPROXIMATE BERGMAN PROJECTIONS

5.1. Preliminaries. In this section we state and prove a result (Th. 5.1 below), which we will use to prove the asymptotic expansion in Th. 2.8 in the next section. The result in question is a modified version of [Berman et al. [6], Prop. 2.5, p. 9]. Our proof is elementary but rather lengthy, and the reader may find it worthwhile to look at the result and go to the next section at the first read.

5.2. A convention. It will be convenient to be able to define integrals $\int_S \chi(w)Y(w)d\mu(w)$ where S is a μ -measurable subset of $\mathbb{C}$, χ a cut-off function, and Y a complex-valued μ -measurable function which is well-defined on $\text{supp } \chi$, but not necessarily on the entire domain S . By convention, we extend the integrand $Y(w)\chi(w)$ to $\mathbb{C}$ by defining $\chi(w)Y(w) = 0$ whenever $\chi(w) = 0$, i.e., we define

$$\int_S \chi(w)Y(w)d\mu(w) := \int_{S \cap \text{supp } \chi} \chi(w)Y(w)d\mu(w).$$

5.3. Statement of the result. Recall the definition of the set $X = \{\Delta Q > 0\}$, as well as of the holomorphic extension ψ of Q from the anti-diagonal, see (2.6). For the approximating kernel K_m^1 , see Definition 2.7. By ∂S we mean the positively oriented boundary of a compact set S . We have the following theorem.

Theorem 5.1. *Let Q be real-analytic in $\mathbb{C}$. Let S be a compact subset of X and fix a point $z_0 \in S$. Then there exists*

- (1) *a number $\varepsilon > 0$ small enough that $K_m^1(z, w)$ makes sense and is Hermitian for all $z, w \in \mathbb{D}(z_0; 2\varepsilon)$ and all $z_0 \in S$,*
- (2) *a function $\chi \in C_0^\infty(\mathbb{C})$ with $\chi = 1$ in $\mathbb{D}(z_0; 3\varepsilon/2)$, $\chi = 0$ outside $\mathbb{D}(z_0; 2\varepsilon)$, and $\|\bar{\partial}\chi\|_{L^2} \leq 3$,*
- (3) *a real-analytic function $v_0(z, w)$ defined for $z, w \in \mathbb{D}(z_0; 2\varepsilon)$,*

such that with

$$(5.1) \quad I_m u(z) = \int_S u(w) \chi(w) K_m^1(z, w) e^{-mQ(w)} dA(w), \quad u \in A_{mQ}^2, \quad z \in \mathbb{D}(z_0; \varepsilon) \cap S^\circ,$$

we have

$$\left| I_m u(z) - u(z) - \frac{1}{2\pi i} \int_{\partial S} u(w) \chi(w) e^{m(\psi(z, \bar{w}) - Q(w))} \left(\frac{1}{z - w} + v_0(z, w) \right) dw \right| \leq \frac{C}{m^{3/2}} e^{mQ(z)/2} \|u\|_{mQ},$$

for all u and z as above, with a number C which only depends on ε and z_0 . Moreover, the numbers C and ε may be chosen independently of z_0 for $z_0 \in S$.

In particular, if $z_0 \in S^\circ$ and if ε is small enough that $\mathbb{D}(z_0; 2\varepsilon) \subset S^\circ$, then $\chi = 0$ on ∂S and so

$$(5.2) \quad |I_m u(z) - u(z)| \leq C m^{-3/2} e^{mQ(z)/2} \|u\|_{mQ}, \quad u \in A_{mQ}^2, \quad z \in \mathbb{D}(z_0; \varepsilon).$$

Remark 5.2. The function v_0 is constructed in the proof, and it depends only on the function ψ and its derivatives. See (5.29) below.

In the case when z_0 is in S° and ε is small enough that $\mathbb{D}(z_0; 2\varepsilon) \subset S^\circ$, the conclusion (5.2) makes it seem reasonable to call the functional $u \mapsto I_m u(z)$ an *approximate Bergman projection*, at least locally, for z in a neighbourhood of z_0 .

5.4. A rough outline; previous work. In the proof we shall construct three operators I_m^j , $j = 1, 2, 3$, such that $I_m u(z) = I_m^1 u(z) + I_m^2 u(z) + I_m^3 u(z)$ for z near z_0 . The constructions will be made so that

$$\begin{aligned} I_m^1 u(z) & \text{ is close to } u(z) + \frac{1}{2\pi i} \int_{\partial S} \frac{u(w) \chi(w) e^{m(\psi(z, \bar{w}) - Q(w))}}{z - w} dw, \\ I_m^2 u(z) & \text{ is close to } \frac{1}{2\pi i} \int_{\partial S} u(w) \chi(w) v_0(z, w) e^{m(\psi(z, \bar{w}) - Q(w))} dw, \\ \text{and } I_m^3 u(z) & \text{ is "negligible" for large } m. \end{aligned}$$

The operator I_m^1 is easy to construct directly, whereas the definitions of the other operators I_m^j for $j = 2, 3$ are somewhat more involved. We have found it convenient to start by considering I_m^1 .

Our approach is based on the paper [6] by Berman et al., in which a somewhat different situation is treated (notably, Q is assumed strictly subharmonic in the entire plane in [6], which means that a somewhat different inner product $|u|_{mQ}^2 = \int |u|^2 e^{-mQ} \Delta Q dA$ is available).

We have found the remarks of Berman in [3], p. 9 quite useful; the construction in [6] is *local* and hence the requirement that Q be globally strictly subharmonic can, at least in principle, be removed. We have found it motivated to give a detailed proof for this statement.

Our approach is heavily inspired by that of the aforementioned papers, but is frequently more elementary. The rest of this section is devoted to the proof of Th. 5.1.

5.5. The first approximation. Throughout the proof, we keep a point $z_0 \in S$ arbitrary but fixed, where S is a given compact subset of X . Also fix a function $u \in A_{mQ}^2$.

The idea is to construct a suitable complex phase function $\phi_z(w)$, such that for fixed z in S° close to z_0 , the main contribution in (5.1) is of the form

$$(5.3) \quad I_m^1 u(z) = \int_S \frac{u(w) \chi(w)}{z - w} \bar{\partial}_w \left(e^{m\phi_z(w)(z-w)} \right) dA(w).$$

To construct ϕ_z , we first define, for points z, w , and $\bar{\zeta}$ sufficiently near each other,

$$(5.4) \quad \theta(z, w, \bar{\zeta}) = \int_0^1 \partial_1 \psi(w + t(z - w), \bar{\zeta}) dt.$$

Then θ is holomorphic in a neighbourhood of the subset $\{(z, z, \bar{z}); z \in \mathbb{C}\} \subset \mathbb{C}^3$ and

$$\theta(z, w, \bar{\zeta})(z - w) = \psi(z, \bar{\zeta}) - \psi(w, \bar{\zeta}).$$

Let $\varepsilon > 0$ be sufficiently small that $\mathbb{D}(z_0; 2\varepsilon) \Subset X$ and such that the functions $\psi(z, w)$, $b_0(z, w)$ and $b_1(z, w)$ make sense and are holomorphic on $\{(z, w); z, \bar{w} \in \mathbb{D}(z_0; 2\varepsilon)\}$ (see Definition 2.7). Writing

$$\delta_0 = \Delta Q(z_0)/2,$$

we may then use (2.9) to choose $\varepsilon > 0$ somewhat smaller if necessary so that

$$(5.5) \quad 2 \operatorname{Re}\{\theta(z, w, \bar{w})(z - w)\} = 2 \operatorname{Re} \psi(z, \bar{w}) - 2Q(w) \leq -\delta_0 |z - w|^2 + Q(z) - Q(w),$$

for all $z, w \in \mathbb{D}(z_0; 2\varepsilon)$. Note that the same ε can be used for all z_0 in K .

Next, we fix a cut-off function $\chi \in C_0^\infty(\mathbb{C})$ such that $\chi = 1$ in $\mathbb{D}(z_0; 3\varepsilon/2)$, $\chi = 0$ outside $\mathbb{D}(z_0; 2\varepsilon)$, $0 \leq \chi \leq 1$ on $\mathbb{C}$ and $\|\bar{\partial}\chi\|_{L^2} \leq 3$, see Subsect. 3.6. Take a point $z \in S^\circ \cap \mathbb{D}(z_0; \varepsilon)$. We now put

$$\phi_z(w) = \theta(z, w, \bar{w}), \quad \text{which means that} \quad \phi_z(w)(z - w) = \psi(z, \bar{w}) - Q(w),$$

and consider the corresponding integral $I_m^1 u(z)$ given by (5.3).

We have the following lemma. The result should be compared with [6], Prop. 2.1.

Lemma 5.3. *There exist positive numbers C_1 and δ depending only on z_0 such that*

$$(5.6) \quad \left| I_m^1 u(z) - \frac{1}{2\pi i} \int_{\partial S} \frac{u(w)\chi(w)e^{m\phi_z(w)(z-w)}}{z-w} dw - u(z) \right| \leq C_1 \varepsilon^{-1} e^{m(Q(z)/2 - \delta\varepsilon^2)} \|u\|_{mQ},$$

for all $u \in A_{mQ}^2$ and all $z \in \mathbb{D}(z_0; \varepsilon) \cap S^\circ$. The numbers C_1 and δ may also be chosen independently of the point $z_0 \in S$. (Indeed, one may take $\delta = \Delta Q(z_0)\varepsilon^2/16$ and $C_1 = 6/\varepsilon$.)

Proof. Keep $z \in \mathbb{D}(z_0; \varepsilon) \cap S^\circ$ arbitrary but fixed and take $u \in A_{mQ}^2$. Since $\chi(z) = 1$ and since $|w - z| \geq \varepsilon/2$ when $\bar{\partial}\chi(w) \neq 0$, (5.3) and Cauchy's formula implies (keep in mind that $z \in S^\circ$)

$$(5.7) \quad \begin{aligned} u(z) + \frac{1}{2\pi i} \int_{\partial S} \frac{u(w)\chi(w)e^{m\phi_z(w)(z-w)}}{z-w} dw &= \int_S \frac{\bar{\partial}_w (u(w)\chi(w)e^{m\phi_z(w)(z-w)})}{z-w} dA(w) = \\ &= I_m^1 u(z) + \int_S \frac{u(w)\bar{\partial}\chi(w)}{z-w} e^{m\phi_z(w)(z-w)} dA(w) = \\ &= I_m^1 u(z) + \int_{\{w \in S; |w-z| \geq \varepsilon/2\}} \frac{u(w)\bar{\partial}\chi(w)}{z-w} e^{m\phi_z(w)(z-w)} dA(w). \end{aligned}$$

Applying the estimate (5.5) to the last integral in (5.7) gives

$$\begin{aligned} \left| \int_{\{w \in S; |w-z| \geq \varepsilon/2\}} \frac{u(w)\bar{\partial}\chi(w)}{z-w} e^{m\phi_z(w)(z-w)} dA(w) \right| &\leq \\ &\leq \int_{\{w \in \mathbb{C}; |w-z| \geq \varepsilon/2\}} \frac{|u(w)\bar{\partial}\chi(w)|}{|z-w|} e^{m(Q(z)/2 - Q(w)/2 - \delta_0 |z-w|^2/2)} dA(w) \leq \\ &\leq (2/\varepsilon) e^{m(Q(z)/2 - \delta_0 \varepsilon^2/8)} \int_{\mathbb{C}} |u(w)\bar{\partial}\chi(w)| e^{-mQ(w)/2} dA(w) \leq \\ &\leq (2/\varepsilon) e^{m(Q(z)/2 - \delta_0 \varepsilon^2/8)} \|u\|_{mQ} \|\bar{\partial}\chi\|_{L^2} \leq (6/\varepsilon) e^{m(Q(z)/2 - \delta_0 \varepsilon^2/8)}, \end{aligned}$$

where we have used that $\|\bar{\partial}\chi\|_{L^2} \leq 3$ to get the last inequality. Recalling that $\delta_0 = \Delta Q(z_0)/2$ and combining with (5.7), we get an estimate

$$\left| u(z) + \frac{1}{2\pi i} \int_{\partial S} \frac{u(w)\chi(w)e^{m\phi_z(w)(z-w)}}{z-w} dw - I_m^1 u(z) \right| \leq (6/\varepsilon) e^{m(Q(z)/2 - \Delta Q(z_0)\varepsilon^2/16)}.$$

The proof is finished. $\square$

5.6. The change of variables. Keep $z_0 \in S$ fixed and note that the definition of θ (see (5.4)) implies that

$$(5.8) \quad [\partial_3 \theta](z, z, \bar{z}) = [\partial_1 \partial_2 \psi](z, \bar{z}) = \Delta Q(z),$$

for all $z \in \mathbb{C}$. Since $\Delta Q(z_0) > 0$, (5.8) implies that there exists a neighbourhood U of $(z, w, \zeta) = (z_0, z_0, \bar{z}_0)$ such that the function

$$F : U \rightarrow F(U) \quad : \quad (z, w, \zeta) \mapsto (z, w, \theta(z, w, \zeta))$$

defines a biholomorphic change of coordinates. In particular, we may regard ζ as a function of the parameters z, w and θ when (z, w, ζ) is in U .

In view of (5.8) we can choose $\varepsilon > 0$ such that $[\partial_3 \theta](z, w, \zeta) \neq 0$ whenever $z, w, \bar{\zeta} \in \mathbb{D}(z_0; 2\varepsilon)$. Since $\phi_z(w) = \theta(z, w, \bar{w})$, we have

$$(5.9) \quad [\partial_3 \theta](z, w, \bar{w}) = \bar{\partial} \phi_z(w),$$

and we infer that when z is fixed in $\mathbb{D}(z_0; \varepsilon)$ we have

$$\bar{\partial} \phi_z(w) \neq 0 \quad \text{when} \quad w \in \mathbb{D}(z_0; 2\varepsilon),$$

Moreover, choosing $\varepsilon > 0$ somewhat smaller if necessary, we can assume that $U = \{(z, w, \zeta); z, w, \bar{\zeta} \in \mathbb{D}(z_0; 2\varepsilon)\}$. Again, note that the same ε can be used for all z_0 in S .

We shall in the following fix a number $\varepsilon > 0$ with the above properties, in addition to the properties which we required earlier, i.e., $\mathbb{D}(z_0; 2\varepsilon) \subseteq X$, the inequality (5.5) holds for all $z, w \in \mathbb{D}(z_0; 2\varepsilon)$, and the functions ψ, b_0 and b_1 are holomorphic in the set $\{(z, w); z, \bar{w} \in \mathbb{D}(z_0; 2\varepsilon)\}$.

5.7. The functions Θ_0 and Υ_0 and a formula for I_m . We now use the correspondence F to define a holomorphic function $\tilde{\psi}$ in $\mathbb{D}(z_0; 2\varepsilon) \times F(U)$ via

$$\tilde{\psi}(z, \xi, w, \theta) = \psi(z, \zeta(\xi, w, \theta)).$$

We also put

$$(5.10) \quad \Theta_0(z, w, \theta) = [\partial_1 \partial_4 \tilde{\psi}](z, z, w, \theta) = b_0(z, \zeta) \cdot [\partial_\theta \zeta](z, w, \theta) = b_0(z, \zeta) / [\partial_\zeta \theta](z, w, \zeta), \quad \zeta = \zeta(z, w, \theta),$$

where we recall that

$$b_0 = \partial_1 \partial_2 \psi.$$

The function Θ_0 was put to use by Berman et al. in [[6], p. 9], where it is called Δ_0 .

Note that (5.4) implies that $\theta(z, z, \zeta) = \partial_1 \psi(z, \zeta)$ for all z and ζ , so that, $[\partial_\zeta \theta](z, z, \zeta) = b_0(z, \zeta)$. Hence (5.10) gives

$$(5.11) \quad \Theta_0(z, z, \theta) = b_0(z, \zeta) / b_0(z, \zeta) = 1.$$

Thus putting

$$\Upsilon_0(z, w, \theta) = - \int_0^1 [\partial_2 \Theta_0](z, z + t(w - z), \theta) dt,$$

we obtain the relation

$$(5.12) \quad \Upsilon_0(z, w, \theta)(z - w) = \Theta_0(z, w, \theta) - 1.$$

We have the following lemma.

Lemma 5.4.

$$(5.13) \quad I_m u(z) = \int_S \frac{u(w) \chi(w)}{z - w} \left(1 + \frac{b_1(z, \bar{w})}{m b_0(z, \bar{w})} \right) \Theta_0(z, w, \phi_z(w)) \bar{\partial}_w \left(e^{m \phi_z(w)(z - w)} \right) dA(w).$$

Proof. Combining (5.10) with (5.9) we get the identity

$$(5.14) \quad \Theta_0(z, w, \phi_z(w)) = b_0(z, \bar{w}) / \bar{\partial} \phi_z(w).$$

Recalling that $K_m^1(z, w) = (mb_0(z, \bar{w}) + b_1(z, \bar{w}))e^{m\psi(z, \bar{w})}$ and $\phi_z(w) = \theta(z, w, \bar{w})$ we thus have (see (5.1))

$$\begin{aligned} I_m u(z) &= \int_S u(w) \chi(w) K_m^1(z, \bar{w}) e^{-mQ(w)} dA(w) = \\ &= \int_S u(w) \chi(w) (mb_0(z, \bar{w}) + b_1(z, \bar{w})) e^{m\phi_z(w)(z-w)} dA(w) = \\ &= \int_S u(w) \chi(w) (mb_0(z, \bar{w}) + b_1(z, \bar{w})) \frac{\bar{\partial}_w (e^{m\phi_z(w)(z-w)})}{m(z-w) \bar{\partial} \phi_z(w)} dA(w) = \\ &= \int_S \frac{u(w) \chi(w)}{z-w} \left(1 + \frac{b_1(z, \bar{w})}{mb_0(z, \bar{w})} \right) \frac{b_0(z, \bar{w})}{\bar{\partial} \phi_z(w)} \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w) = \\ &= \int_S \frac{u(w) \chi(w)}{z-w} \left(1 + \frac{b_1(z, \bar{w})}{mb_0(z, \bar{w})} \right) \Theta_0(z, w, \phi_z(w)) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w), \end{aligned}$$

where we have used (5.14) to get the last identity. $\square$

Comparing (5.13) with the formula (5.3), we now see a connection between $I_m u(z)$ and $I_m^1 u(z)$. We shall exploit this relation within short.

5.8. The differential equation. We will now show that the functions b_0 , b_1 and Θ_0 are connected via an important differential equation.

Lemma 5.5. *For all $z, \bar{\zeta} \in \mathbb{D}(z_0; 2\varepsilon)$, we have the identity*

$$(5.15) \quad \frac{b_1(z, \zeta)}{b_0(z, \zeta)} = - \left(\frac{\partial^2}{\partial w \partial \theta} \Theta_0(z, w, \theta) \right) \Big|_{z=w, \theta=\theta(z, z, \zeta)}.$$

Proof. In view of (2.7), it suffices to show that the holomorphic function

$$(5.16) \quad B_1(z, \zeta) = -b_0(z, \zeta) \cdot [\partial_2 \partial_3 \Theta_0](z, z, \theta(z, z, \zeta))$$

satisfies $B_1(z, \zeta) = \frac{1}{2} \frac{\partial}{\partial \bar{\zeta}} \left(\frac{1}{b_0(z, \bar{\zeta})} \frac{\partial}{\partial z} b_0(z, \zeta) \right)$ for all z and $\bar{\zeta}$ in a neighbourhood of z_0 .

To this end, we first note that a Taylor expansion yields that

$$\psi(w, \zeta) = \psi(z, \zeta) + (w-z) \frac{\partial \psi}{\partial z}(z, \zeta) + \frac{(w-z)^2}{2} \frac{\partial^2 \psi}{\partial z^2}(z, \zeta) + O((w-z)^3), \quad \text{as } w \rightarrow z,$$

i.e.,

$$\theta(z, w, \zeta) = \frac{\psi(w, \zeta) - \psi(z, \zeta)}{w-z} = \frac{\partial \psi}{\partial z}(z, \zeta) + \frac{w-z}{2} \frac{\partial^2 \psi}{\partial z^2}(z, \zeta) + O((w-z)^2).$$

Differentiating with respect to ζ and using that $b_0 = \partial_1 \partial_2 \psi$ yields

$$(5.17) \quad \frac{\partial \theta}{\partial \zeta} = b_0(z, \zeta) + \frac{w-z}{2} \frac{\partial b_0}{\partial z}(z, \zeta) + O((w-z)^2).$$

Dividing both sides of (5.17) by $b_0 = b_0(z, \zeta)$ and using (5.10), we get

$$\frac{1}{\Theta_0} = \frac{1}{b_0} \frac{\partial \theta}{\partial \zeta} = 1 + \frac{w-z}{2b_0} \frac{\partial b_0}{\partial z} + O((w-z)^2).$$

Inverting this relation we obtain

$$\Theta_0 = b_0 / \frac{\partial \theta}{\partial \zeta} = 1 - \frac{w-z}{2b_0} \frac{\partial b_0}{\partial z} + O((w-z)^2).$$

In view of (5.16), it yields that

$$\begin{aligned}
B_1(z, \zeta(z, z, \theta)) &= -b_0(z, \zeta) \cdot \frac{\partial^2}{\partial \theta \partial w} \Theta_0(z, w, \theta) \Big|_{w=z} = \\
&= \frac{b_0(z, \zeta)}{2} \frac{\partial}{\partial \theta} \left(\frac{1}{b_0(z, \zeta)} \frac{\partial}{\partial z} b_0(z, \zeta) \right) \Big|_{z=w} = \\
&= \frac{b_0(z, \zeta)}{2} \frac{\partial}{\partial \zeta} \left(\frac{1}{b_0(z, \zeta)} \frac{\partial}{\partial z} b_0(z, \zeta) \right) \Big|_{z=w} / \frac{\partial \theta}{\partial \zeta}(z, z, \zeta) = \\
&= \frac{1}{2} \frac{\partial}{\partial \zeta} \left(\frac{1}{b_0(z, \zeta)} \frac{\partial}{\partial z} b_0(z, \zeta) \right) = b_1(z, \zeta).
\end{aligned}$$

Here we have used (5.10), (5.11) and (2.7) to obtain the last equality. $\square$

5.9. The operator ∇_m . For the further analysis, it will be convenient to consider a differential operator ∇_m defined for complex smooth functions A of the parameters z, w and θ by

$$(5.18) \quad \nabla_m A = \frac{\partial A}{\partial \theta} + m(z - w)A = e^{-m\theta(z-w)} \frac{\partial}{\partial \theta} (e^{m\theta(z-w)} A(z, w, \theta)).$$

Cf. [6], p. 5. It will be useful to note that, evaluating at a point where $\theta = \phi_z(w)$ we have

$$(5.19) \quad \nabla_m A \Big|_{\theta=\phi_z(w)} = \frac{e^{-m\phi_z(w)(z-w)}}{\bar{\partial}\phi_z(w)} \bar{\partial}_w (e^{m\phi_z(w)(z-w)} A(z, w, \phi_z(w))),$$

as is easily verified by carrying out the differentiation and comparing with (5.18). From this, we now derive an important identity,

$$\begin{aligned}
(5.20) \quad & \int_S u(w) \chi(w) [\nabla_m A](z, w, \phi_z(w)) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w) = \\
&= m \int_S u(w) \chi(w) \bar{\partial}_w (e^{m\phi_z(w)(z-w)} A(z, w, \phi_z(w))) dA(w) = \\
&= -m \int_S u(w) \bar{\partial} \chi(w) e^{m\phi_z(w)(z-w)} A(z, w, \phi_z(w)) dA(w) + \\
&\quad + \frac{m}{2\pi i} \int_{\partial S} u(w) \chi(w) e^{m\phi_z(w)(z-w)} A(z, w, \phi_z(w)) dw = \\
&= -m \int_{\{w \in S; |z-w| \geq \varepsilon/2\}} u(w) \bar{\partial} \chi(w) e^{m\phi_z(w)(z-w)} A(z, w, \phi_z(w)) dA(w) + \\
&\quad + \frac{m}{2\pi i} \int_{\partial S} u(w) \chi(w) e^{m\phi_z(w)(z-w)} A(z, w, \phi_z(w)) dw, \quad z \in \mathbb{D}(z_0; \varepsilon).
\end{aligned}$$

Here we have used (5.19) to deduce the first equality. The second equality follows from Green's formula. To get the last equality we have used that $|z - w| \geq \varepsilon/2$ when $\bar{\partial} \chi(w) \neq 0$.

We now have the following lemma, which is based on [[6], Lemma 2.3 and the discussion on p. 9].

Lemma 5.6. *There exist holomorphic functions $A_m(z, w, \theta)$ and $B(z, w, \theta)$, uniformly bounded on $F(U)$, such that*

$$(5.21) \quad \left(1 + \frac{b_1(z, \zeta)}{mb_0(z, \zeta)} \right) \Theta_0(z, w, \theta) = 1 + m^{-1} \nabla_m A_m(z, w, \theta) + m^{-2} B(z, w, \theta),$$

where it is understood that $\theta = \theta(z, w, \zeta)$. Moreover, we have that

$$(5.22) \quad \lim_{m \rightarrow \infty} A_m = \Upsilon_0,$$

with uniform convergence on $F(U)$.

Proof. Put

$$\Theta_1(z, w, \theta) = \frac{b_1(z, \zeta)}{b_0(z, \zeta)} \Theta_0(z, w, \theta),$$

so that the left hand side in (5.21) can be written

$$(5.23) \quad \left(1 + \frac{b_1(z, \zeta)}{mb_0(z, \zeta)}\right) \Theta_0(z, w, \theta) = \Theta_0(z, w, \theta) + m^{-1} \Theta_1(z, w, \theta).$$

Next, recall that $\Theta_0(z, w, \theta) - 1 = (z - w) \Upsilon_0(z, w, \theta)$, cf. (5.12). Further, from the relation (5.15), which reads

$$\frac{b_1(z, \zeta)}{b_0(z, \zeta)} = - \left(\frac{\partial^2}{\partial \theta \partial w} \Theta_0(z, w, \theta) \right) \Big|_{z=w}, \quad \zeta = \zeta(z, z, \theta),$$

it follows that

$$(5.24) \quad \Theta_1(z, z, \theta) = - \left(\frac{\partial^2}{\partial \theta \partial w} \Theta_0(z, w, \theta) \right) \Big|_{z=w} \cdot \Theta_0(z, z, \theta) = - \left(\frac{\partial^2}{\partial \theta \partial w} \Theta_0(z, w, \theta) \right) \Big|_{z=w},$$

where we have used that $\Theta_0(z, z, \theta) = 1$ (see (5.11)) to get the last equality. (5.24) allows us to write

$$\Theta_1(z, w, \theta) + \partial_\theta \partial_w \Theta_0(z, w, \theta) = (z - w) \Upsilon_1(z, w, \theta),$$

where Υ_1 is holomorphic. Now define

$$A_m = \Upsilon_0 + m^{-1}(\Upsilon_1 - (\partial_\theta \partial_w \Upsilon_0)) \quad , \quad B = -\partial_\theta(\Upsilon_1 - (\partial_\theta \partial_w \Upsilon_0)).$$

Then A_m and B are holomorphic and $A_m \rightarrow \Upsilon_0$ as $m \rightarrow \infty$.

We will see that straightforward calculations show that

$$(5.25) \quad \nabla_m A_m = m(\Theta_0 - 1) + \Theta_1 + m^{-1} \frac{\partial}{\partial \theta} \left(\frac{\Theta_1}{z - w} - \frac{\partial_\theta \Theta_0}{(z - w)^2} \right) = m(\Theta_0 - 1) + \Theta_1 - m^{-1} B.$$

Indeed, when $z \neq w$ we have

$$\begin{aligned} A_m &= \frac{\Theta_0 - 1}{z - w} + m^{-1} \left(\frac{\Theta_1 + \partial_\theta \partial_w \Theta_0}{z - w} - \partial_w \left(\frac{\partial_\theta \Theta_0}{z - w} \right) \right) = \\ &= \frac{\Theta_0 - 1}{z - w} + m^{-1} \left(\frac{\Theta_1}{z - w} - \frac{\partial_\theta \Theta_0}{(z - w)^2} \right) \end{aligned}$$

so that

$$\partial_\theta A_m = \frac{\partial_\theta \Theta_0}{z - w} + m^{-1} \left(\frac{\partial_\theta \Theta_1}{z - w} - \frac{\partial_\theta^2 \Theta_0}{(z - w)^2} \right),$$

and

$$m(z - w)A_m = m(\Theta_0 - 1) + \Theta_1 - \frac{\partial_\theta \Theta_0}{z - w},$$

which gives (5.25), since $\nabla_m A_m = \partial_\theta A_m + m(z - w)A_m$.

Dividing through by m in the (5.25) now gives

$$\Theta_0 + m^{-1} \Theta_1 = 1 + m^{-1} \nabla_m A_m + m^{-2} B,$$

and glancing at (5.23) we obtain (5.21). □

5.10. **Decomposition of I_m .** In view of (5.13) and Lemma 5.6, we can now write

$$\begin{aligned}
 I_m u(z) &= \int_S \frac{u(w)\chi(w)}{z-w} \left(1 + \frac{b_1(z, \bar{w})}{mb_0(z, \bar{w})} \right) \Theta_0(z, w, \bar{w}) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w) = \\
 &= \int_S \frac{u(w)\chi(w)}{z-w} \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w) + \\
 (5.26) \quad &+ \frac{1}{m} \int_S \frac{u(w)\chi(w)}{z-w} [\nabla_m A_m](z, w, \phi_z(w)) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w) + \\
 &+ \frac{1}{m^2} \int_S \frac{u(w)\chi(w)}{z-w} B(z, w, \phi_z(w)) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w),
 \end{aligned}$$

where A_m and B are uniformly bounded and holomorphic near $(z_0, z_0, \phi_{z_0}(z_0))$.

We recognize the first integral in the right hand side of (5.26) as $I_m^1 u(z)$ (see (5.3)). Let us denote the others by

$$I_m^2 u(z) = \frac{1}{m} \int_S \frac{u(w)\chi(w)}{z-w} \nabla_m A_m(z, w, \phi_z(w)) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w),$$

and

$$(5.27) \quad I_m^3 u(z) = \frac{1}{m^2} \int_S \frac{u(w)\chi(w)}{z-w} B(z, w, \phi_z(w)) \bar{\partial}_w (e^{m\phi_z(w)(z-w)}) dA(w).$$

5.11. **Estimates for I_m^2 .** We start by estimating $I_m^2 u(z)$ when $z \in \mathbb{D}(z_0; \varepsilon)$, and note that (5.20) yields that

$$\begin{aligned}
 I_m^2 u(z) &= - \int_{\{w \in S; |z-w| \geq \varepsilon/2\}} u(w) \bar{\partial} \chi(w) e^{m\phi_z(w)(z-w)} A_m(z, w, \phi_z(w)) dA(w) + \\
 (5.28) \quad &+ \frac{1}{2\pi i} \int_{\partial S} u(w) \chi(w) e^{m\phi_z(w)(z-w)} A_m(z, w, \phi_z(w)) dw.
 \end{aligned}$$

Let us now define

$$(5.29) \quad v_0(z, w) = \Upsilon_0(z, w, \phi_z(w)),$$

so that $A_m(z, w, \phi_z(w)) \rightarrow v_0(z, w)$ with uniform convergence for $z, w \in \mathbb{D}(z_0; 2\varepsilon)$ when $m \rightarrow \infty$, by Lemma 5.6. Also let $C' = C'(\varepsilon)$ be an upper bound for $\{|A_m(z, w, \phi_z(w))|; z, w \in \mathbb{D}(z_0; 2\varepsilon), m \geq 1\}$. We now use (5.28) and the estimate (5.5) to get

$$\begin{aligned}
 &\left| I_m^2 u(z) - \frac{1}{2\pi i} \int_{\partial S} u(w) \chi(w) e^{m\phi_z(w)(z-w)} v_0(z, w) dw \right| \leq \\
 (5.30) \quad &\leq C' \int_{\{w \in S; |w-z| \geq \varepsilon/2\}} |u(w) \bar{\partial} \chi(w)| e^{m(Q(z)/2 - Q(w)/2 - \delta_0 |z-w|^2/2)} dA(w) \leq \\
 &\leq C' e^{m(Q(z)/2 - \delta_0 \varepsilon^2/8)} \int_{\mathbb{C}} |u(w) \bar{\partial} \chi(w)| e^{-Q(w)/2} dA(w) \leq \\
 &\leq C' e^{m(Q(z)/2 - \delta_0 \varepsilon^2/8)} \|u\|_{mQ} \|\bar{\partial} \chi\|_{L^2} \leq C_2 e^{m(Q(z)/2 - \delta \varepsilon^2)} \|u\|_{mQ},
 \end{aligned}$$

where $\delta = \delta_0/8 = \Delta Q(z_0)/16$, and $C_2 = 3C'$.

5.12. **Estimates for I_m^3 .** To estimate $I_m^3 u(z)$, we note that (5.27) implies

$$I_m^3 u(z) = \frac{1}{m} \int_{\mathbb{C}} u(w) \chi(w) B(z, w, \phi_z(w)) e^{m\phi_z(w)(z-w)} \bar{\partial} \phi_z(w) dA(w).$$

It suffices to estimate this integral using (5.5). This gives

$$|I_m^3 u(z)| \leq \frac{1}{m} \int_{\mathbb{C}} |u(w) \chi(w) B(z, w, \phi_z(w))| e^{m(Q(z)/2 - Q(w)/2 - m\delta_0 |z-w|^2/2)} |\bar{\partial} \phi_z(w)| dA(w).$$

Since the function $|B(z, w, \phi_z(w))\bar{\partial}\phi_z(w)\chi(w)|$ can be estimated by a number C_3 independent of z and w for $z \in \mathbb{D}(z_0; \varepsilon)$ and $w \in \mathbb{D}(z_0; 2\varepsilon)$, it gives, in view of the Cauchy–Schwarz inequality

$$(5.31) \quad \begin{aligned} |I_m^3 u(z)| &\leq C_3 m^{-1} e^{mQ(z)/2} \int_{\mathbb{C}} |u(w)\chi(w)| e^{-mQ(w)/2} dA(w) \leq \\ &\leq C_3 m^{-1} e^{mQ(z)/2} \|u\|_{mQ} \left(\int_{\mathbb{C}} e^{-m\delta_0|z-w|^2} dA(w) \right)^{1/2} \leq C_3 \delta_0^{-1/2} m^{-3/2} e^{mQ(z)/2} \|u\|_{mQ}, \end{aligned}$$

where $C_3 = C_3(\varepsilon) = \sup \{ |B(z, w, \phi_z(w))\bar{\partial}\phi_z(w)\chi(w)|; z, w \in \mathbb{D}(z_0; 2\varepsilon) \}$.

5.13. Conclusion of the proof of Theorem 5.1. By (5.26) and the estimates (5.6), (5.30) and (5.31), (using that $\phi_z(w)(z-w) = \psi(z, \bar{w}) - Q(w)$)

$$\begin{aligned} &\left| I_m u(z) - u(z) - \frac{1}{2\pi i} \int_{\partial S} u(w)\chi(w) \left(\frac{1}{z-w} + \nu_0(z, w) \right) e^{m(\psi(z, \bar{w}) - Q(w))} dw \right| \leq \\ &\leq \left| I_m^1 u(z) - u(z) - \frac{1}{2\pi i} \int_{\partial S} \frac{u(w)\chi(w)e^{m\phi_z(w)(z-w)}}{z-w} dw \right| + \\ &\quad + \left| I_m^2 u(z) - \frac{1}{2\pi i} \int_{\partial S} u(w)\chi(w)\nu_0(z, w)e^{m\phi_z(w)(z-w)} dw \right| + |I_m^3 u(z)| \leq \\ &\leq C_1 \varepsilon^{-1} e^{m(Q(z)/2 - \delta\varepsilon^2)} \|u\|_{mQ} + C_2 e^{m(Q(z)/2 - \delta\varepsilon^2)} \|u\|_{mQ} + C_3 \delta_0^{-1} m^{-3/2} e^{mQ(z)/2} \|u\|_{mQ}, \quad z \in \mathbb{D}(z_0; \varepsilon) \cap S^\circ. \end{aligned}$$

It thus suffices to choose a number C large enough that

$$(5.32) \quad C_1 e^{-\delta m} + C_2 e^{-\delta m} + C_3 m^{-3/2} \leq C m^{-3/2}, \quad m \geq 1.$$

Now recall that the numbers C_1 , C_2 , and C_3 only depend on ε , where $\varepsilon > 0$ can be chosen independently of z_0 for z_0 in the given compact subset S of X . Moreover, we have that $\delta = \Delta Q(z_0)\varepsilon^2/16$, and $\Delta Q(z_0)$ is bounded below by a positive number for $z_0 \in S$. It follows that C in (5.32) can be chosen independently of $z_0 \in S$, and the proof is finished. $\square$

6. THE PROOF OF THEOREM 2.8

6.1. Preliminaries. In this section we prove Th. 2.8. Our approach which follows [[6], Sect. 3] is obtained by assembling the information from Lemma 3.2, Th. 5.1, and Cor. 4.5. To facilitate the presentation, we divide the proof into steps.

First recall that adding a constant to Q means that $K_{m,n}$ is only changed by a multiplicative constant, and hence we can (and will) assume that $Q \geq 1$ on $\mathbb{C}$.

Fix a compact subset $K \Subset S_\tau^\circ \cap X$ and a point $z_0 \in S_\tau^\circ \cap X$, and let $\varepsilon > 0$ and $\chi \in C_0^\infty(\mathbb{C})$ be as in Th. 5.1. Choosing $\varepsilon > 0$ somewhat smaller if necessary, we may ensure that $2\varepsilon < \text{dist}(K, \mathbb{C} \setminus (S_\tau \cap X))$ and that

$$(6.1) \quad |K_m^1(z, w)|^2 e^{-m(Q(z)+Q(w))} \leq C m^2 e^{-\delta_0 m|z-w|^2}, \quad z, \bar{w} \in \mathbb{D}(z_0; 2\varepsilon), \quad z_0 \in K,$$

for some positive numbers C and δ_0 (see (2.11)). We also introduce the set

$$\Sigma = \{z \in \mathbb{C}; \text{dist}(z, K) \leq 2\varepsilon\} \Subset S_\tau^\circ \cap X.$$

Our goal is to prove an estimate

$$|K_{m,n}(z, w) - K_m^1(z, w)| e^{-m(Q(z)+Q(w))/2} \leq C m^{-1}, \quad z, w \in \mathbb{D}(z_0; \varepsilon), \quad n \geq m\tau - M, \quad m \geq m_0,$$

where $M \geq 0$ is given, and C and m_0 are independent of the specific choice of the point $z_0 \in K$.

In the following, we let C denote various (more or less) absolute constants which can change meaning from time to time, even within the same chain of inequalities.

Let $P_{m,n} : L_{mQ}^2 \rightarrow H_{m,n}$ denote the orthogonal projection, that is,

$$P_{m,n}u(z) = \int_{\mathbb{C}} u(w)K_{m,n}(z, w)e^{-mQ(w)}dA(w).$$

Throughout the proof we fix two points $z, w \in \mathbb{D}(z_0; \varepsilon)$ and introduce the functions

$$(6.2) \quad u_z(\zeta) = K_{m,n}(\zeta, z) \quad \text{and} \quad v_w(\zeta) = \begin{cases} \chi(\zeta)K_m^1(\zeta, w), & \zeta \in \mathbb{D}(z_0; 2\varepsilon), \\ 0, & \text{otherwise,} \end{cases}$$

as well as

$$(6.3) \quad d_w(\zeta) = v_w(\zeta) - P_{m,n}v_w(\zeta), \quad \zeta \in \mathbb{C}.$$

We note that $v_w \in C_0^\infty(\mathbb{C})$ with $\text{supp } v_w \subset \Sigma \Subset \mathcal{S}_\tau^\circ \cap X$.

Lemma 6.1. *There exists a positive number C independent of $m \geq 1$, n , z and w , such that*

$$(6.4) \quad |K_{m,n}(z, w) - P_{m,n}v_w(z)| \leq Cm^{-1}e^{m(Q(z)+Q(w))/2}.$$

Moreover, C can be chosen independent of z_0 for $z_0 \in K$.

Proof. By Th. 5.1, there is a number C depending only on ε and K , such that

$$|I_m u_z(w) - u_z(w)| \leq Cm^{-3/2}e^{mQ(w)/2}\|u_z\|_{mQ},$$

where I_m is given by (5.1) with $S = \Sigma$. Next note that the estimate (3.2) implies

$$\|u_z\|_{mQ}^2 = K_{m,n}(z, z) \leq Cme^{mQ(z)},$$

with a number C depending only on τ . We conclude that

$$(6.5) \quad |I_m u_z(w) - K_{m,n}(w, z)| \leq Cm^{-1}e^{m(Q(z)+Q(w))/2}.$$

We next note that

$$(6.6) \quad \overline{I_m u_z(w)} = \int_{\mathbb{C}} \chi(\zeta)K_m^1(\zeta, w)K_{m,n}(z, \zeta)e^{-mQ(\zeta)}dA(\zeta) = P_{m,n}v_w(z),$$

with v_w given by (6.2). Thus the statement (6.4) is immediate from (6.6) and (6.5). $\square$

Lemma 6.2. *Let two positive numbers M_0 and M_1 be given such that (4.17) is fulfilled, and put $m_0 = \max\{(1+M_0)/\tau, 2M_0\}$. Then for all m, n such that $n \geq (m-M_0)\tau + M_1$ and $m \geq m_0$, there exist positive numbers δ and C , independent of z_0, m, n, z , and w , such that*

$$(6.7) \quad |d_w(z)|^2 \leq Cm^3e^{-m\delta}e^{m(Q(z)+Q(w))}.$$

Proof. The function d_w is the $L_{mQ,n}^2$ -minimal solution to the $\bar{\partial}$ -equation $\bar{\partial}d_w = \bar{\partial}v_w$. Since $\text{supp } v_w \subset \mathcal{S}_\tau$, Cor. 4.5 yields that

$$(6.8) \quad \|d_w\|_{mQ}^2 \leq C\|\bar{\partial}v_w\|_{mQ}^2, \quad m \geq m_0, \quad n \geq (m-M_0)\tau + M_1,$$

with a number C depending only on τ, M_0 and M_1 . But $\bar{\partial}v_w(\zeta) = \bar{\partial}\chi(\zeta)K_m^1(\zeta, w)$, whence the estimate (6.1) shows that there are numbers C and $\delta_0 > 0$ such that

$$(6.9) \quad |\bar{\partial}v_w(\zeta)|^2 e^{-m(Q(\zeta)+Q(w))} \leq Cm^2|\bar{\partial}\chi(\zeta)|^2 e^{-m\delta_0|\zeta-w|^2}, \quad \zeta \in \mathbb{C}.$$

Now, since $|\zeta - w| \geq \varepsilon/2$ whenever $\bar{\partial}\chi(\zeta) \neq 0$, we deduce from (6.9) that

$$(6.10) \quad |\bar{\partial}v_w(\zeta)|^2 e^{-mQ(\zeta)} \leq Cm^2|\bar{\partial}\chi(\zeta)|^2 e^{m(Q(w)-\delta)}, \quad \zeta \in \mathbb{C},$$

where $\delta = \delta_0\varepsilon^2/4$. Using (6.8) and then integrating the inequality (6.10) with respect to $dA(\zeta)$, we get

$$(6.11) \quad \|d_w\|_{mQ}^2 \leq C\|\bar{\partial}v_w\|_{mQ}^2 \leq Cm^2e^{m(Q(w)-\delta)}\|\bar{\partial}\chi\|_{L^2}^2 \leq Cm^2e^{m(Q(w)-\delta)}.$$

Next note that the function d_w is holomorphic in the disk $\mathbb{D}(z_0; 3\varepsilon/2)$, so that Lemma 3.2 gives

$$(6.12) \quad |d_w(z)|^2 e^{-mQ(z)} \leq Cm \int_{\mathbb{D}(z; \varepsilon/\sqrt{m})} |d_w(\zeta)|^2 e^{-mQ(\zeta)} dA(\zeta) \leq Cm \|d_w\|_{mQ}^2,$$

where C only depends on ε and τ . Combining (6.12) with (6.11), we end up with (6.7), and so we are done. $\square$

6.2. Conclusion of the proof of Theorem 2.8. Since $\chi(z) = 1$, we have that $v_w(z) = K_m^1(z, w)$, whence by (6.3)

$$|K_{m,n}(z, w) - K_m^1(z, w)| = |K_{m,n}(z, w) - v_w(z)| \leq |K_{m,n}(z, w) - P_{m,n}v_w(z)| + |d_w(z)|.$$

In view of Lemmas 6.1 and 6.2, the right hand side can be estimated by

$$C(m^{-1} + m^{3/2}e^{-m\delta/2})e^{m(Q(z)+Q(w))/2},$$

whenever $n \geq (m\tau - M_0) + M_1$ and $m \geq m_0$. For $m \geq 1$, the latter expression is dominated by $Cm^{-1}e^{m(Q(z)+Q(w))/2}$. The proof is finished. $\square$

7. BEREZIN QUANTIZATION AND GAUSSIAN CONVERGENCE

7.1. Preliminaries. In this section we use the expansion formula for $K_{m,n}$ (Th. 2.8) to prove theorems 2.3 and 2.6. In the proofs, we set $\tau = 1$ and $m = n$. (The argument in the general case follows the same pattern.) It then becomes natural to write K_n for $K_{n,n}$ etc. We also fix a compact subset $K \Subset \mathcal{S}_1^\circ \cap X$, a point $z_0 \in K$, and a positive number ε with the properties listed in Th. 2.8, and we put

$$\delta_n = \varepsilon \log n / \sqrt{n}.$$

7.2. The proof of Theorem 2.3. It suffices to show that

$$(7.1) \quad B_n^{\langle z_0 \rangle}(\mathbb{D}(z_0; \delta_n)) \rightarrow 1 \quad \text{as } n \rightarrow \infty.$$

Indeed, since $B_n^{\langle z_0 \rangle}$ is a p.m., this implies Th. 2.3.

In order to prove (7.1), we apply Th. 2.8, which gives

$$K_n(z_0, z)e^{-n(Q(z)+Q(z_0))/2} = (nb_0(z_0, \bar{z}) + b_1(z_0, \bar{z}))e^{n(\operatorname{Re} \psi(z_0, \bar{z}) - (Q(z)+Q(z_0))/2)} + O(n^{-1}), \quad z \in \mathbb{D}(z_0; \varepsilon)$$

when $n \rightarrow \infty$, where the O is uniform for $z_0 \in K$. In view of (2.9), we have

$$(7.2) \quad K_n(z_0, z)e^{-n(Q(z)+Q(z_0))/2} = (nb_0(z_0, \bar{z}) + O(1))e^{n(-\Delta Q(z_0)/2 + R(z_0, z))} + O(n^{-1}), \quad z \in \mathbb{D}(z_0; \varepsilon)$$

with a function R satisfying $|R(z, w)| \leq C|z - w|^3$ whenever $z, w \in \mathbb{D}(z_0; \varepsilon)$ and $z_0 \in K$. Note that

$$(7.3) \quad n|R(z_0, z)| \leq Cn\delta_n^3 \leq C\log^3 n / \sqrt{n}, \quad \text{when } z \in \mathbb{D}(z_0; \delta_n), \quad z_0 \in K.$$

Introducing this estimate in (7.2) gives

$$|K_n(z_0, z)|^2 e^{-n(Q(z)+Q(z_0))} = (n^2 |b_0(z_0, \bar{z})|^2 + O(n))e^{-n\Delta Q(z_0)|z_0 - z|^2 + O(\log^3 n / \sqrt{n})} + O(n^{-2}),$$

for $z \in \mathbb{D}(z_0; \delta_n)$ as $n \rightarrow \infty$, where the O -terms are uniform for $z_0 \in K$. Furthermore, (2.8) yields that

$$(7.4) \quad \frac{|K_n(z_0, z)|^2}{K_n(z_0, z_0)} e^{-nQ(z)} = \frac{|K_n(z_0, z)|^2}{n\Delta Q(z_0) + O(1)} e^{-n(Q(z_0)+Q(z))}, \quad z \in \mathbb{D}(z_0; \varepsilon),$$

as $n \rightarrow \infty$. Note that the left hand side in (7.4) is the density $B_n^{\langle z_0 \rangle}(z)$, so that (7.2) and (7.4) implies

$$(7.5) \quad B_n^{\langle z_0 \rangle}(z) = \frac{n^2 |b_0(z_0, \bar{z})|^2 + O(n)}{n\Delta Q(z_0) + O(1)} e^{-n\Delta Q(z_0)|z - z_0|^2 + O(\log^3 n / \sqrt{n})} + O(n^{-2}), \quad z \in \mathbb{D}(z_0; \delta_n).$$

Integrating (7.5) with respect to dA over $\mathbb{D}(z_0; \delta_n)$ and using the mean-value theorem for integrals now gives that there are positive numbers v_{n,z_0} converging to 1 (uniformly for $z_0 \in K$), and also

complex numbers b_{n,z_0} converging to $b_0(z_0, \bar{z}_0) = \Delta Q(z_0)$ (uniformly for $z_0 \in K$) as $n \rightarrow \infty$, such that

$$\begin{aligned} B_n^{\langle z_0 \rangle}(\mathbb{D}(z_0; \delta_n)) &= v_{n,z_0} \frac{n^2 |b_{n,z_0}|^2 + O(n)}{n\Delta Q(z_0) + O(1)} \int_{\mathbb{D}(z_0; \delta_n)} e^{-n\Delta Q(z_0)|z-z_0|^2} dA(z) + O(n^{-2}) \int_{\mathbb{D}(z_0; \delta_n)} dA(z) = \\ &= v_{n,z_0} \frac{n^2 |b_{n,z_0}|^2 + O(n)}{n^2 \Delta Q(z_0)^2 + O(n)} (1 - e^{-\Delta Q(z_0)\varepsilon^2 \log^2 n}) + O(n^{-2}). \end{aligned}$$

The expression in the right hand side converges to 1 as $n \rightarrow \infty$. This proves (7.1), and Th. 2.3 follows. $\square$

7.3. The proof of Theorem 2.6. It suffices to show that there are numbers ε_n converging to zero as $n \rightarrow \infty$ such that

$$(7.6) \quad \int_{\mathbb{C}} \left| B_n^{\langle z_0 \rangle}(z) - n\Delta Q(z_0)e^{-n\Delta Q(z_0)|z-z_0|^2} \right| dA(z) \leq \varepsilon_n, \quad z_0 \in K.$$

Indeed, (2.5) follows from (7.6) after the change of variables $z \mapsto z_0 + z/\sqrt{n\Delta Q(z_0)}$ in the integral in (7.6).

To prove (7.6), we split the integral with respect to the decomposition $\{|z - z_0| < \delta_n\} \cup \{|z - z_0| \geq \delta_n\}$, and put

$$\begin{aligned} A_{n,z_0} &= \int_{\{|z-z_0| < \delta_n\}} \left| B_n^{\langle z_0 \rangle}(z) - n\Delta Q(z_0)e^{-n\Delta Q(z_0)|z-z_0|^2} \right| dA(z), \\ B_{n,z_0} &= \int_{\{|z-z_0| \geq \delta_n\}} \left| B_n^{\langle z_0 \rangle}(z) - n\Delta Q(z_0)e^{-n\Delta Q(z_0)|z-z_0|^2} \right| dA(z). \end{aligned}$$

Considering A_{n,z_0} first, we get from (7.6) that

$$A_{n,z_0} = \int_{\mathbb{D}(z_0; \delta_n)} \left| e^{nR(z_0, \bar{z})} \frac{n^2 |b_0(z_0, \bar{z})|^2 + O(n)}{n\Delta Q(z_0) + O(1)} - n\Delta Q(z_0) \right| e^{-n\Delta Q(z_0)|z-z_0|^2} dA(z) + O(n^{-2}),$$

as $n \rightarrow \infty$. Next we put

$$s_{n,z_0} = \sup_{z \in \mathbb{D}(z_0; \delta_n)} \left\{ \left| e^{nR(z_0, \bar{z})} \frac{n^2 |b_0(z_0, \bar{z})|^2 + O(n)}{n\Delta Q(z_0) + O(1)} - n\Delta Q(z_0) \right| \right\},$$

and observe (7.3) implies that $s_{n,z_0}/n \rightarrow 0$, uniformly for $z_0 \in K$. It yields that

$$A_{n,z_0} \leq s_{n,z_0} \int_{\mathbb{D}(z_0; \delta_n)} e^{-n\Delta Q(z_0)|z-z_0|^2} dA(z) + O(n^{-2}) \leq C s_{n,z_0}/n,$$

which converges to 0 as $n \rightarrow \infty$ uniformly for $z_0 \in K$.

To estimate B_{n,z_0} we simply observe that

$$(7.7) \quad B_{n,z_0} \leq \int_{\{|z-z_0| \geq \delta_n\}} B_n^{\langle z_0 \rangle}(z) dA(z) + \int_{\{|z-z_0| \geq \delta_n\}} n\Delta Q(z_0)e^{-n\Delta Q(z_0)|z-z_0|^2} dA(z).$$

Since $B_n^{\langle z_0 \rangle}$ is a p.m., the estimate (7.1) yields that the first integral in the right hand side of (7.7) converges to 0 as $n \rightarrow \infty$. Moreover, a simple calculation yields that the second integral in (7.7) converges to 0 when $n \rightarrow \infty$. We have shown that $B_{n,z_0} \rightarrow 0$ as $n \rightarrow \infty$ with uniform convergence for $z_0 \in K$. The proof is finished. $\square$

8. OFF-DIAGONAL DAMPING

8.1. An estimate for $K_{m,n}$. In this section we prove Th. 2.4. We shall obtain that theorem from Th. 8.3 below, which is of independent interest, and has applications in random matrix theory, see [1]. Our analysis depends on the following lemma. It will be convenient to define the set

$$\mathcal{S}_{\tau,1} = \{\zeta; \text{dist}(\zeta, \mathcal{S}_\tau) \leq 1\}.$$

Lemma 8.1. *Assume that $Q \in C^2(\mathbb{C})$. Let $z_0 \in \mathcal{S}_\tau \cap X$ and let M be given non-negative numbers. Put*

$$8d = \text{dist}(z_0, \mathbb{C} \setminus (\mathcal{S}_\tau \cap X)) \quad a = \inf\{\Delta Q(\zeta); \zeta \in \mathbb{D}(z_0; 6d)\} \quad A = \sup\{\Delta Q(\zeta); \zeta \in \mathcal{S}_{\tau,1}\}.$$

There then exists positive numbers C and ϵ such that for all $m, n \geq 1$ with $m\tau - M \leq n \leq m\tau + 1$, we have

$$(8.1) \quad |K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq Cm^2 e^{-\epsilon \sqrt{m} \min\{d, |z_0-z|\}}, \quad z \in \mathcal{S}_\tau.$$

Here C depends only on M, a, A , and τ , while ϵ only depends on a, τ and M .

Remark 8.2. For fixed τ and M , our method of proof gives that ϵ can be chosen proportional to a while C can be chosen proportional to a^{-1} . Indeed, our proof shows that if there is a positive number c depending only on τ and M such that, with $a' = \min\{a, 1\}$,

$$|K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq C \frac{m^3}{am+c} e^{-\epsilon' a' \sqrt{m} \min\{d, |z_0-z|\}}, \quad z \in \mathcal{S}_\tau,$$

with C and ϵ' independent of a . This estimate can easily be extended to all $z \in \mathbb{C}$ by adapting the proof of Th. 8.3 below.

Before we turn to the proof of Lemma 8.1, we use it to prove the main result of this section.

Theorem 8.3. *In the situation of Lemma 8.1 we also have an estimate*

$$(8.2) \quad |K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq Cm^2 e^{-\epsilon \sqrt{m} \min\{d, |z_0-z|\}} e^{-m(Q(z)-\widehat{Q}_\tau(z))}, \quad z \in \mathbb{C},$$

valid for all $z_0 \in \mathcal{S}_\tau \cap X$, $m \geq 1$, and all n with $m\tau - M \leq n \leq m\tau + 1$.

Proof. In view of Lemma 8.1, and since $Q = \widehat{Q}_\tau$ on $\mathcal{S}_\tau$, it suffices to show the estimate (8.2) when $z \notin \mathcal{S}_\tau$. To this end, we consider the function

$$f(\zeta) = \log |K_{m,n}(z_0, \zeta)|^2 - m\widehat{Q}_\tau(\zeta).$$

Since $\widehat{Q}_\tau$ is harmonic on $\mathbb{C} \setminus \mathcal{S}_\tau$, f is subharmonic there. Moreover, since $n-1 \leq m\tau$ and since $K_{m,n}(\cdot, z_0) \in H_{m,n}$, we have a simple estimate

$$\log |K_{m,n}(z_0, \zeta)|^2 \leq (n-1) \log |\zeta|^2 + O(1) \leq m\tau \log |\zeta|^2 + O(1) \quad \text{when } \zeta \rightarrow \infty,$$

while the relation (2.3) says that $\widehat{Q}_\tau(\zeta) = \tau \log |\zeta|^2 + O(1)$ when $\zeta \rightarrow \infty$. Hence f is bounded above. Furthermore, it is clear that f is harmonic in a punctured neighbourhood of ∞ , which yields that f has a representation $f(\zeta) = h(\zeta) - c \log |\zeta|$ for all large enough $|\zeta|$, where c is a non-negative number and h is harmonic at ∞ (see e.g. [[21], Cor. 0.3.7, p. 12]). In particular, f extends to a subharmonic function on $\mathbb{C}^* \setminus \mathcal{S}_\tau$. Finally, since Lemma 8.1 yields that

$$f(\zeta) \leq \log(Cm^2) + mQ(z_0) - \epsilon \sqrt{m}d, \quad \text{when } \zeta \in \partial \mathcal{S}_\tau,$$

the maximum principle shows that the same estimate holds for all $\zeta \in \mathbb{C}^* \setminus \mathcal{S}_\tau$. This means that

$$(8.3) \quad |K_{m,n}(z_0, z)|^2 e^{-m\widehat{Q}_\tau(z)} \leq Cm^2 e^{-\epsilon \sqrt{m}d} e^{mQ(z_0)}, \quad \text{when } z \notin \mathcal{S}_\tau,$$

which implies (8.2) when $z \notin \mathcal{S}_\tau$. □

Background. Estimates related to those considered in Lemma 8.1 are known, see e.g. Lindholm's article [20], Prop. 9, pp. 404–407. We will follow the basic strategy used in that paper in our proof below. Since we are considering a different situation with polynomial Bergman kernels (instead of the full Bergman kernel of A^2_{mQ}), and since we are not assuming the weight Q to be globally strictly subharmonic, nontrivial modifications of the classical arguments are needed. Our main tool for accomplishing this is provided by the weighted L^2 estimates in Th. 4.4.

8.2. The proof of Lemma 8.1. We can and will assume that $Q \geq 1$ on $\mathbb{C}$. Moreover, we will denote various constants by the same letter C , which can change meaning during the course of the calculations. When C depends on one or several parameters, we will always specify this. To simplify the proof we will first reduce the problem by treating three simple cases.

8.3. Case 1: $md^2 \leq 1$. Since we have assumed that $n \leq m\tau + 1$, the estimate (3.3) of Prop. 3.6 applies. It gives that

$$(8.4) \quad |K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq Cm^2, \quad z \in \mathbb{C},$$

with a number C depending only on τ . The assertion (8.1) follows immediately in case $d = 0$. In the remaining case we have $d > 0$ and $m \leq d^{-2}$. Then, for any $\epsilon > 0$ and any $z \in \mathbb{C}$, we have that $e^{-\epsilon \sqrt{m} \min\{d, |z_0 - z|\}} \geq e^{-\epsilon d^{-1}d} = e^{-\epsilon}$, so that

$$(8.5) \quad |K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq Ce^\epsilon e^{-\epsilon \sqrt{m} \min\{d, |z_0 - z|\}}.$$

This gives the desired estimate (8.1) with C replaced by Ce^ϵ .

8.4. Notation. We now fix positive numbers M_0 and M_1 such that the relation (4.17) is satisfied. We further choose M_0 and M_1 so that $[(m - M_0)\tau + M_1] \leq m\tau - M$ for all $m \geq 1$, and let $m_0 = \max\{(1 + M_0)/\tau, 4M_0\}$. Let n be a positive integer such that $n \leq m\tau + 1$. We also fix a point $z \in \mathcal{S}_\tau$ and let R be a number such that $\mathcal{S}_\tau \subset \mathbb{D}(0; R)$.

8.5. Case 2: $m \leq m_0$. Given any $\epsilon > 0$, we have that $e^{-\epsilon \sqrt{m} \min\{|z_0 - z|, d\}} \geq e^{-\epsilon \sqrt{m_0} R}$, and so (8.4) implies that

$$|K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq Ce^\epsilon \sqrt{m_0} R e^{-\epsilon \sqrt{m} \min\{|z_0 - z|, d\}}.$$

Thus (8.1) holds with C replaced by $Ce^\epsilon \sqrt{m_0} R$.

8.6. Case 3: $|z - z_0| \leq 8/\sqrt{m}$. In this case, $e^{-\epsilon \sqrt{m} \min\{d, |z_0 - z|\}} \geq e^{-8\epsilon}$, and thus (8.4) implies

$$|K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0)+Q(z))} \leq Ce^{8\epsilon} m^2 e^{-\epsilon \sqrt{m} \min\{d, |z_0 - z|\}}.$$

We have shown (8.1) in the case when $|z_0 - z| \leq 8/\sqrt{m}$ with C replaced by $Ce^{8\epsilon}$.

8.7. Case 4: $m \geq m_0$, $md^2 \geq 1$ and $|z - z_0| \geq 8/\sqrt{m}$. In the sequel we fix any integer n with $n \geq [(m - M_0)\tau + M_1]$. Here $[(m - M_0)\tau + M_1] > 0$ for all $m \geq m_0$ by our choice of m_0 (see Subsect. 8.4).

It is important to note that the assumption $md^2 \geq 1$ means that $1/\sqrt{m} \leq d$ so that

$$8/\sqrt{m} \leq 8d = \text{dist}(z_0, \mathbb{C} \setminus (\mathcal{S}_\tau \cap X)).$$

Our starting point is Lemma 3.3, which yields that

$$(8.6) \quad |K_{m,n}(z, z_0)|^2 e^{-mQ(z)} \leq Cm \int_{\mathbb{D}(z; 1/\sqrt{m})} |K_{m,n}(\zeta, z_0)|^2 e^{-mQ(\zeta)} dA(\zeta), \quad z \in \mathcal{S}_\tau,$$

where the number C only depends on A . Define

$$\varepsilon_0(\zeta) = \min\{|z_0 - \zeta|/2, 4d\}, \quad \zeta \in \mathbb{C}.$$

Note that

$$(8.7) \quad 4/\sqrt{m} \leq \varepsilon_0(z) \leq 4d.$$

Let χ_0 be a smooth non-negative function such that

$$(8.8) \quad \chi_0 = 0 \quad \text{on} \quad \mathbb{D}(z_0; \varepsilon_0(z)/2) \quad \text{and} \quad \chi_0 = 1 \quad \text{outside} \quad \mathbb{D}(z_0; \varepsilon_0(z)),$$

and also $|\bar{\partial}\chi_0|^2 \leq (C/\varepsilon_0(z)^2)\chi_0$ with C an absolute constant ($C = 5$ will do). In view of (8.7), it yields that

$$|\bar{\partial}\chi_0|^2 \leq Cm\chi_0 \quad \text{on} \quad \mathbb{C},$$

where $C = 5/16$. Notice that $\bar{\partial}\chi_0$ is supported in the annulus

$$(8.9) \quad U_0 = U_0(z_0, z) = \{\zeta; \varepsilon_0(z)/2 \leq |z_0 - \zeta| \leq \varepsilon_0(z)\}.$$

Since χ_0 is non-negative on $\mathbb{C}$ and since $\chi_0 = 1$ on $\mathbb{D}(z; 1/\sqrt{m})$, the estimate (8.6) implies that

$$(8.10) \quad |K_{m,n}(z, z_0)|^2 e^{-mQ(z)} \leq Cm \int_{\mathbb{C}} \chi_0(\zeta) |K_{m,n}(\zeta, z_0)|^2 e^{-mQ(\zeta)} dA(\zeta),$$

where C only depends on A . Let $H_{\chi_0, mQ}$ be the linear space $H_{m,n}$ with inner product

$$\langle f, g \rangle_{\chi_0, mQ} = \int_{\mathbb{C}} f \bar{g} \chi_0 e^{-mQ} dA.$$

We rewrite integral in the the right hand side in (8.10) in the following way

$$(8.11) \quad \begin{aligned} \int_{\mathbb{C}} \chi_0(\zeta) |K_{m,n}(\zeta, z_0)|^2 e^{-mQ(\zeta)} dA(\zeta) &= \sup \left\{ |\langle u, K_{m,n}(\cdot, z_0) \rangle_{\chi_0, mQ}|^2; u \in H_{m,n}, \int_{\mathbb{C}} |u|^2 \chi_0 e^{-mQ} dA \leq 1 \right\} = \\ &= \sup \left\{ \left| \langle \chi_0 u, K_{m,n}(\cdot, z_0) \rangle_{mQ} \right|^2; u \in H_{m,n}, \int_{\mathbb{C}} |u|^2 \chi_0 e^{-mQ} dA \leq 1 \right\} = \\ &= \sup \left\{ |P_{m,n}[\chi_0 u](z_0)|^2; u \in H_{m,n}, \int_{\mathbb{C}} |u|^2 \chi_0 e^{-mQ} dA \leq 1 \right\}, \end{aligned}$$

where $P_{m,n}$ is the orthogonal projection of L^2_{mQ} onto $H_{m,n}$. Now fix $u \in H_{m,n}$ with $\int_{\mathbb{C}} |u|^2 \chi_0 e^{-mQ} dA \leq 1$ and recall that $P_{m,n}[\chi_0 u] = \chi_0 u - u_*$, where u_* is the $L^2_{mQ, n}$ -minimal solution to $\bar{\partial}u_* = \bar{\partial}(\chi_0 u) = u \bar{\partial}\chi_0$. In particular, u_* is holomorphic in $\mathbb{D}(z_0; \varepsilon_0(z)/2)$ and $u_* = -P_{m,n}[\chi_0 u]$ there. See Subsect. 4.1.

We intend to apply Th. 4.4 with a suitable real-valued function ϱ_m . We shall at first only specify ϱ_m by requiring certain properties of it. An explicit construction of ϱ_m is then given at the end of the proof.

Condition 1. We require that

$$(8.12) \quad \varrho_m = 0 \quad \text{on} \quad \mathbb{D}(z_0; 1/(2\sqrt{m})),$$

Condition 2. There exists a number $\epsilon > 0$ depending only on a, M_0 and τ , such that

$$\varrho_m(\zeta) \leq -\epsilon \sqrt{m} \varepsilon_0(z)/4 \quad \text{when} \quad |\zeta - z_0| \geq \varepsilon_0(z)/2,$$

Condition 3. The various conditions on ϱ_m in Th. 4.4 are satisfied. More precisely,

(i) ϱ_m is $C^{1,1}$ -smooth and

$$(m - M_0)\Delta Q(\zeta) + \Delta \varrho_m(\zeta) \geq ma/2, \quad \text{for a.e.} \quad \zeta \in \overline{\mathbb{D}}(z_0; 6d),$$

(ii) ϱ_m is constant in $\mathbb{C} \setminus \overline{\mathbb{D}}(z_0; 6d)$,

(iii) We have that

$$\frac{|\bar{\partial}\varrho_m|^2}{ma/2} \leq \frac{1}{4e^{M_0 q_\tau}} \quad \text{on} \quad \mathbb{C},$$

where $q_\tau = \sup_{S_\tau} \{Q(\zeta)\}$.

It is clear that (i), (ii) and (iii) imply the conditions on ϱ_m in Th. 4.4, with $\kappa = 1/2$.

We now turn to consequences of the conditions on ϱ_m , and start with condition 1. Applying Lemma 3.2, we find that, for any real function ϱ_m satisfying (8.12), we have

$$(8.13) \quad |u_*(z_0)|^2 e^{-mQ(z_0)} \leq Cm \int_{\mathbb{D}(z_0; 1/(2\sqrt{m}))} |u_*(\zeta)|^2 e^{-mQ(\zeta)} dA(\zeta) \leq Cm \int_{\mathbb{C}} |u_*(\zeta)|^2 e^{\varrho_m(\zeta) - mQ(\zeta)} dA(\zeta),$$

where C depends only on A .

By Condition 3, we may apply Th. 4.4 to the integral in the right hand side in (8.13). It yields that

$$(8.14) \quad \int_{\mathbb{C}} |u_*|^2 e^{\varrho_m - mQ} dA \leq C \int_{U_0} |u \bar{\partial} \chi_0|^2 \frac{e^{\varrho_m - mQ}}{ma/2 + M_1 c_\tau} dA,$$

with a number C depending only on τ and M_0 (note that $(1-\kappa)^{-2} = 4$). Here $c_\tau = \inf_{S_\tau} \{(1+|\zeta|^2)^{-2}\} \geq (1+R^2)^{-2}$ and U_0 is the annulus defined in (8.9).

But since $|\bar{\partial} \chi_0|^2 \leq Cm \chi_0$, (8.14) implies

$$\int_{\mathbb{C}} |u_*|^2 e^{\varrho_m - mQ} dA \leq \frac{Cm}{am + M_1 c_\tau} \int_{U_0} |u|^2 \chi_0 e^{\varrho_m - mQ} dA.$$

Here C only depends on M_0 and τ . We now use Condition 2, which implies that $\varrho_m(\zeta) \leq -\epsilon \sqrt{m} \varepsilon_0(z)$ whenever $\zeta \in U_0$. It yields that we may continue to estimate

$$(8.15) \quad \int_{U_0} |u|^2 \chi_0 e^{\varrho_m - mQ} dA \leq e^{-\epsilon \sqrt{m} \varepsilon_0(z)/4} \int_{\mathbb{C}} |u|^2 \chi_0 e^{-mQ} dA \leq e^{-\epsilon \sqrt{m} \min\{|z_0 - z|/8, d\}},$$

where we have used that $\int_{\mathbb{C}} |u|^2 \chi_0 e^{-mQ} dA \leq 1$ in the last step.

Tracing back through (8.10)–(8.15), we infer that

$$|K_{m,n}(z_0, z)|^2 e^{-m(Q(z_0) + Q(z))} \leq C \frac{m^3}{am + M_1 c_\tau} e^{-\epsilon \sqrt{m} \min\{|z_0 - z|/8, d\}} \leq Ca^{-1} m^2 e^{-\epsilon \sqrt{m} \min\{|z_0 - z|/8, d\}},$$

where C depends on M_0 , τ , and A . This proves Lemma 8.1 (with $\epsilon/8$ instead of ϵ and Ca^{-1} in place of C) under the hypotheses that a function ϱ_m satisfying conditions 1, 2, and 3 above exists. To finish the proof we must verify the existence of such a ϱ_m .

8.8. Construction of ϱ_m . We now construct a function ϱ_m and a positive number ϵ which satisfy the conditions 1, 2, and 3 above. We look for a radial function of the form

$$\varrho_m(\zeta) = -\epsilon \sqrt{m} S_m(|\zeta - z_0|),$$

where the number $\epsilon > 0$ will be fixed later. We start by giving an explicit construction of S_m , the proof of conditions 1 through 3 will then be accomplished without difficulty.

We recall that $1/\sqrt{m} \leq d$ and start by specifying the derivative S'_m to be the piecewise linear continuous function on $[0, \infty)$ such that

$$S'_m = 0 \quad \text{on} \quad [0, 1/(2\sqrt{m})] \cup [6d, \infty) \quad \text{and} \quad S'_m = 1 \quad \text{on} \quad [1/\sqrt{m}, 5d],$$

and S'_m is affine on each of the intervals $[1/(2\sqrt{m}), 1/\sqrt{m}]$ and $[5d, 6d]$. The distributional derivative of S'_m is then a linear combination of characteristic functions,

$$S''_m = 2\sqrt{m} \mathbf{1}_{[1/(2\sqrt{m}), 1/\sqrt{m}]} - (1/d) \mathbf{1}_{[5d, 6d]},$$

so that (since $md^2 \geq 1$)

$$|S''_m| \leq \max\{2\sqrt{m}, 1/d\} = 2\sqrt{m}.$$

We now define S_m by requiring that $S_m(0) = 0$. Since $S'_m = 0$ on $[0, 1/(2\sqrt{m})]$ it is then clear that $S_m = 0$ on $[0, 1/(2\sqrt{m})]$. Moreover, when $2/\sqrt{m} \leq t \leq 5d$, we get

$$S_m(t) = \int_0^t S'_m(s) ds \geq t - 1/\sqrt{m} \geq t/2, \quad t \in [2/\sqrt{m}, 5d],$$

since $S'_m = 1$ on $[1/\sqrt{m}, 5d]$. When $t \geq 5d$, we plainly have

$$S_m(t) \geq 5d - 1/\sqrt{m} \geq 4d.$$

We conclude that

$$(8.16) \quad S_m(t) \geq \min\{t/2, 4d\}, \quad t \geq 2/\sqrt{m}.$$

In particular, denoting by c_m the constant value that S_m assumes on $[6d, \infty)$, we have $c_m \geq 4d$. This finishes the construction of S_m , and the corresponding function ϱ_m is clearly of class $C^{1,1}(\mathbb{C})$.

We now verify the conditions 1 through 3 above. First, Condition 1 is clear, since $S_m(t) = 0$ when $t \leq 1/(2\sqrt{m})$. Also, part (ii) of condition 3 is clear; since $S_m(t) = c_m$ is constant when $t \geq 6d$, we have that $\varrho_m(\zeta) = -\epsilon\sqrt{m}c_m$ is constant when $\zeta \notin \overline{\mathbb{D}}(0; 6d)$.

Since $\varepsilon_0(z) \geq 4/\sqrt{m}$, (8.16) implies that $\varrho_m(\zeta) = -\epsilon S_m(|\zeta - z_0|) \leq -\epsilon\varepsilon_0(\zeta)$ when $|\zeta - z_0| \geq \varepsilon_0(z)/2$. Since moreover $\varepsilon_0(\zeta) \geq \varepsilon_0(z)/4$ in this case, we get that condition 2 is satisfied.

There remains to check parts (i) and (iii) of condition 3, and to make precise what we mean by "ε". To this end, we need the following estimates

$$(8.17) \quad \left| \bar{\partial} \varrho_m(\zeta) \right|^2 = \epsilon^2 m \left| S'_m(|\zeta - z_0|) \right|^2 / 4 \leq \epsilon^2 m / 4, \quad \zeta \in \mathbb{C},$$

and

$$(8.18) \quad \left| \Delta \varrho_m(\zeta) \right| \leq \frac{\epsilon}{4} \sqrt{m} \left(\left| S''_m(|\zeta - z_0|) \right| + \frac{S'_m(|z_0 - \zeta|)}{|z_0 - \zeta|} \right) \leq \epsilon m, \quad \zeta \in \mathbb{C},$$

which follows immediately from the properties of S_m (since $|z_0 - \zeta| \geq 1/(2\sqrt{m})$ when $S'_m(|z_0 - \zeta|) \neq 0$).

To verify (i), we use (8.18). It yields that it suffices to choose $\epsilon > 0$ such that $(m - M_0)a - \epsilon m \geq ma/2$ for $m \geq m_0$, i.e. $\epsilon \leq (1/2 - M_0/m)a$. Since we have assumed that $m_0 \geq 4M_0$, it thus suffices to choose $\epsilon = a/4$.

We finally verify (iii). By (8.17), it suffices to choose an $\epsilon > 0$ such that $(\epsilon^2/4)/(a/2) \leq 1/(4e^{M_0q_\tau})$, i.e. $\epsilon^2 \leq a/(2e^{M_0q_\tau})$.

We have verified the existence of $\epsilon > 0$, of the form $\epsilon = \min\{\sqrt{a/(2e^{M_0q_\tau})}, a/2\}$. This shows that the choice $\epsilon = c \min\{a, 1\}$ works with a proportionality constant c which depends on M_0 and q_τ . The proof is finished. $\square$

8.9. The proof of Theorem 2.4. Let K be a compact subset of $\mathcal{S}_\tau^\circ \cap X$, and pick $M \geq 0$. By Th. 2.8 we have that $K_{m,n}(z_0, z_0)e^{-mQ(z_0)} = m\Delta Q(z_0) + O(1)$ as $m \rightarrow \infty$ and $n \geq m\tau - M$, where the O is uniform for $z_0 \in K$. It yields that

$$(8.19) \quad \mathbb{B}_{m,n}^{(z_0)}(z) = \frac{|K_{m,n}(z, z_0)|^2}{K_{m,n}(z_0, z_0)} e^{-mQ(z)} = \frac{|K_{m,n}(z, z_0)|^2}{m\Delta Q(z_0) + O(1)} e^{-m(Q(z) + Q(z_0))}, \quad z \in \mathbb{C},$$

as $m \rightarrow \infty$ and $n \geq m\tau - M$. Since ΔQ is bounded below by a positive number on K , the right hand side in (8.19) can be estimated by

$$Cm^{-1} |K_{m,n}(z, z_0)|^2 e^{-m(Q(z) + Q(z_0))}, \quad z \in \mathbb{C}$$

where C depends on the lower bound of ΔQ on K . The assertion now follows from Th. 8.3. $\square$

9. THE BARGMANN–FOCK CASE AND HARMONIC MEASURE

9.1. Preliminaries. In this section we prove Th. 2.11. We therefore put $Q(z) = |z|^2$. Recall that in this case $\mathcal{S}_\tau = \overline{\mathbb{D}}(0; \sqrt{\tau})$, and (see (2.12))

$$(9.1) \quad dB_{m,n}^{(z_0)}(z) = m \frac{|E_{n-1}(mz\bar{z}_0)|^2}{E_{n-1}(m|z_0|^2)} e^{-m|z|^2} dA(z) \quad \text{where} \quad E_k(z) = \sum_{j=0}^k \frac{z^j}{j!}.$$

9.2. The action on polynomials.

Proposition 9.1. *Fix a complex number $z_0 \neq 0$, a positive integer d and let n be an integer, $n \geq d + 1$. Then, for all analytic polynomials u of degree at most d , we have*

$$(9.2) \quad \text{p. v.} \int_{\mathbb{C}} u(z^{-1}) dB_{m,n}^{(z_0)} \rightarrow u(z_0^{-1}), \quad \text{as } m \rightarrow \infty,$$

uniformly in n , $n \geq d + 1$.

Proof. It is sufficient to prove the statement for $u(z) = z^j$ with $j \leq d$. The left hand side in (9.2) can then be written

$$\text{p. v.} \int_{\mathbb{C}} z^{-j} dB_{m,n}^{(z_0)} = \frac{mb_{m,n}^j(z_0)}{E_{n-1}(m|z_0|^2)},$$

where we have put

$$b_{m,n}^j(z_0) = \text{p. v.} \int_{\mathbb{C}} z^{-j} \left| \sum_{k=0}^{n-1} \frac{(mz_0\bar{z})^k}{k!} \right|^2 e^{-m|z|^2} dA(z).$$

Expanding the square yields

$$b_{m,n}^j(z_0) = \sum_{k,l=0}^{n-1} \frac{m^{k+l} z_0^k \bar{z}_0^l}{k!l!} \text{p. v.} \int_{\mathbb{C}} \bar{z}^k z^{l-j} e^{-m|z|^2} dA(z).$$

Clearly only those k, l for which $k = l - j$ give a non-zero contribution to the sum, and therefore,

$$b_{m,n}^j(z_0) = z_0^{-j} \sum_{l=j}^{n-1} \frac{m^{2l-j} |z_0|^{2l}}{(l-j)!l!} \int_{\mathbb{C}} |z|^{2(l-j)} e^{-m|z|^2} dA(z) = \frac{1}{mz_0^j} \sum_{l=j}^{n-1} \frac{m^l |z_0|^{2l}}{l!}.$$

It follows that

$$b_{m,n}^j(z_0) = \frac{1}{mz_0^j} \sum_{l=j}^{n-1} \frac{(m|z_0|^2)^l}{l!} = \frac{1}{mz_0^j} (E_{n-1}(m|z_0|^2) - E_{j-1}(m|z_0|^2)),$$

and so

$$\frac{mb_{m,n}^j(z_0)}{E_{n-1}(m|z_0|^2)} = \frac{1}{z_0^j} \left(1 - \frac{E_{j-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} \right).$$

Finally, since $j \leq d < n$,

$$\frac{E_{j-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} \leq \frac{E_{d-1}(m|z_0|^2)}{E_d(m|z_0|^2)} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

□

Proposition 9.2. *Let $0 < r < \sqrt{\tau}$, $z_0 \in \mathbb{C} \setminus \overline{\mathbb{D}}(0; \sqrt{\tau})$ and u an analytic polynomial. Then*

$$\text{p. v.} \int_{\mathbb{D}(0;r)} u(z^{-1}) dB_{m,n}^{(z_0)}(z) \rightarrow 0 \quad \text{as } m \rightarrow \infty \quad \text{and } n/m \rightarrow \tau.$$

Proof. Put, for $v = 0, 1, 2, \dots$,

$$b_{m,n}^v(z_0, r) = \text{p. v.} \int_{\mathbb{D}(0;r)} z^{-v} dB_{m,n}^{(z_0)}.$$

A straightforward calculation based on (9.1) leads to

$$(9.3) \quad b_{m,n}^v(z_0, r) = \frac{1}{z_0^v E_{n-1}(m|z_0|^2)} \sum_{j=v}^{n-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^{mr^2} s^{j-v} e^{-s} ds.$$

We suppose n is greater than v by at least two units, so that we may pick an integer k with $v < k < n$, and split the sum (9.3) accordingly:

$$(9.4) \quad b_{m,n}^v(z_0, r) = \frac{1}{z_0^v E_{n-1}(m|z_0|^2)} \left\{ \sum_{j=v}^{k-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^{mr^2} s^{j-v} e^{-s} ds + \sum_{j=k}^{n-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^{mr^2} s^{j-v} e^{-s} ds \right\}.$$

We estimate the first term trivially as follows:

$$(9.5) \quad \sum_{j=v}^{k-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^{mr^2} s^{j-v} e^{-s} ds \leq \sum_{j=v}^{k-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^\infty s^{j-v} e^{-s} ds = \sum_{j=0}^{k-1} \frac{(m|z_0|^2)^j}{j!} = E_{k-1}(m|z_0|^2).$$

As for the second term, we use the fact that the function $s \mapsto s^{j-v} e^{-s}$ is increasing on the interval $[0, j-v]$, to say that

$$\int_0^{mr^2} s^{j-v} e^{-s} ds \leq (mr^2)^{j-v+1} e^{-mr^2},$$

provided that $j \geq mr^2 + v$. It follows that if $k \geq mr^2 + v$, then

$$\sum_{j=k}^{n-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^{mr^2} s^{j-v} e^{-s} ds \leq (mr^2)^{1-v} e^{-mr^2} \sum_{j=k}^{n-1} \frac{(mr|z_0|)^{2j}}{j!(j-v)!}.$$

By Stirling's formula, $j! \geq \sqrt{2\pi} j^{j+1/2} e^{-j}$, so that

$$\frac{(mr|z_0|)^{2j}}{j!} e^{-mr^2} \leq \frac{1}{\sqrt{2\pi} j} m^j |z_0|^{2j} \left(\frac{mr^2}{j} e^{1-\frac{mr^2}{j}} \right)^j.$$

Since the function $x \mapsto x e^{1-x}$ is increasing on the interval $[0, 1]$, it yields that

$$\frac{(mr|z_0|)^{2j}}{j!} e^{-mr^2} \leq \frac{1}{\sqrt{2\pi} j} m^j |z_0|^{2j} \left(\frac{mr^2}{k} e^{1-\frac{mr^2}{k}} \right)^j, \quad mr^2 + v \leq k \leq j.$$

We write

$$(9.6) \quad c_{k,m} = \frac{mr^2}{k} e^{1-\frac{mr^2}{k}} \leq 1, \quad mr^2 + v \leq k,$$

and conclude that

$$(9.7) \quad \begin{aligned} \sum_{j=k}^{n-1} \frac{(m|z_0|^2)^j}{j!(j-v)!} \int_0^{mr^2} s^{j-v} e^{-s} ds &\leq (mr^2)^{1-v} \sum_{j=k}^{n-1} \frac{(m|z_0|^2 c_{k,m})^j}{(j-v)! \sqrt{2\pi} j} \\ &\leq \frac{(mr^2)^{1-v}}{\sqrt{2\pi}} (c_{k,m})^k \sum_{j=k}^{n-1} \frac{(m|z_0|^2)^j}{(j-v)!} \leq \frac{mr^2}{\sqrt{2\pi}} \left(\frac{|z_0|}{r} \right)^{2v} (c_{k,m})^k E_{n-v-1}(m|z_0|^2). \end{aligned}$$

Now, a combination (9.5) and (9.7) applied to (9.4) yields

$$(9.8) \quad \begin{aligned} |z_0^v b_{m,n}^v(z_0, r)| &\leq \frac{E_{k-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} + \frac{mr^2}{\sqrt{2\pi}} \left(\frac{|z_0|}{r} \right)^{2v} (c_{k,m})^k \frac{E_{n-v-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} \\ &\leq \frac{E_{k-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} + \frac{mr^2}{\sqrt{2\pi}} \left(\frac{|z_0|}{r} \right)^{2v} (c_{k,m})^k. \end{aligned}$$

We would like to show that each of the terms on the right hand side of (9.8) can be made small by choosing k cleverly. As for the first term, we appeal to a theorem of Szegő, [23], Hilfssatz 1, p. 54, which states that

$$E_l(lx) = \frac{1}{\sqrt{2\pi} l} (ex)^l \frac{x}{x-1} (1 + \varepsilon_l(x)) \quad x > 1,$$

where $\varepsilon_l(x) \rightarrow 0$ uniformly on compact subintervals of $(1, \infty)$ as $l \rightarrow \infty$. It follows that

$$\frac{E_{k-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} = \sqrt{\frac{n-1}{k-1}} \frac{m|z_0|^2 - n + 1}{m|z_0|^2 - k + 1} \left(\frac{em|z_0|^2}{k-1}\right)^{k-1} \left(\frac{em|z_0|^2}{n-1}\right)^{1-n} \frac{1 + \varepsilon_{k-1}(\frac{m|z_0|^2}{k-1})}{1 + \varepsilon_{n-1}(\frac{m|z_0|^2}{n-1})}.$$

Finally, we decide to pick k such that

$$k/m \rightarrow \beta$$

as $k, m \rightarrow \infty$, where $r^2 < \beta < \tau$. We observe that with this choice of k , the above epsilons tend to zero as $k, m, n \rightarrow \infty$. The function

$$y \mapsto (e/y)^y, \quad 0 < y \leq 1$$

is strictly increasing, so that with

$$y_1 = \frac{k-1}{m|z_0|^2} \approx \frac{\beta}{|z_0|^2}, \quad y_2 = \frac{n-1}{m|z_0|^2} \approx \frac{\tau}{|z_0|^2},$$

we have

$$\frac{(e/y_1)^{y_1}}{(e/y_2)^{y_2}} \leq \theta < 1,$$

where at least for large k, m, n , the number θ may be taken to be independent of k, m, n . It follows that

$$\left(\frac{em|z_0|^2}{k-1}\right)^{k-1} \left(\frac{em|z_0|^2}{n-1}\right)^{1-n} \leq \theta^{-m|z_0|^2},$$

so that

$$\frac{E_{k-1}(m|z_0|^2)}{E_{n-1}(m|z_0|^2)} \leq (1 + o(1)) \sqrt{\frac{\tau}{\beta}} \frac{|z_0|^2 - \tau}{|z_0|^2 - \beta} \theta^{-m|z_0|^2} \rightarrow 0$$

exponentially quickly as $k, m, n \rightarrow \infty$.

Finally, as for the second term, we observe that the numbers $c_{k,m}$ defined by (9.6) have the property that

$$c_{k,m} \rightarrow \frac{r^2}{\beta} e^{1-\frac{r^2}{\beta}} < 1,$$

as $k, m, n \rightarrow \infty$ in the given fashion. In particular, the second term converges exponentially quickly to 0. The proof is complete. $\square$

Corollary 9.3. *Let $z_0 \in \mathbb{C} \setminus \overline{\mathbb{D}}(0; \sqrt{\tau})$, and let ω be an open set in $\mathbb{C}$ which contains the circle $\mathbb{T}(0, \sqrt{\tau})$. Further, let u be an analytic polynomial. Then*

$$\int_{\omega} u(z^{-1}) dB_{m,n}^{(z_0)}(z) \rightarrow u(z_0^{-1}) \quad \text{as } m \rightarrow \infty \text{ and } n/m \rightarrow \tau.$$

Proof. This follows from propositions 9.2 and 2.2. $\square$

9.3. The proof of Theorem 2.11. Let $\mathcal{H}_\tau$ be the class of continuous functions $\mathbb{C}^* \rightarrow \mathbb{C}$ which are harmonic on $\mathbb{C}^* \setminus \mathcal{S}_\tau$. For a function $f \in C_b(\mathbb{C})$ we write $\tilde{f}$ for the unique function of class $\mathcal{H}_\tau$ which coincides with f on $\mathbb{D}(0, \sqrt{\tau})$. We must show that

$$\int_{\mathbb{C}} f(z) dB_{m,n}^{(z_0)}(z) \rightarrow \tilde{f}(z_0)$$

as $m \rightarrow \infty$ and $n/m \rightarrow \tau$. See e.g. [14], p. 90.

Convolving with the Féjer kernel, we see that f may be uniformly approximated by functions which on a neighborhood ω of $\mathbb{T}(0, \sqrt{\tau})$ are of the form $u(z^{-1})$, with u a harmonic polynomial. We may therefore w.l.o.g. suppose that f itself is of this form, i.e. $f(z) = u(z^{-1})$ when $z \in \omega$. Thus $f(z) = \tilde{f}(z) = u(z^{-1})$ on ω . By Prop. 2.2,

$$\int_{\mathbb{C}} (f(z) - \tilde{f}(z)) dB_{m,n}^{(z_0)}(z) \rightarrow 0,$$

as $m \rightarrow \infty$ and $n/m \rightarrow \tau$. Moreover, Cor. 9.3 gives that

$$\int_{\mathbb{C}} \widetilde{f}(z) dB_{m,n}^{\langle z_0 \rangle}(z) = \int_{\mathbb{C}} u(z^{-1}) dB_{m,n}^{\langle z_0 \rangle}(z) \rightarrow u(z_0^{-1}) = \widetilde{f}(z_0),$$

as $m \rightarrow \infty$ and $n/m \rightarrow \tau$. □

REFERENCES

- [1] Ameur, Y., Hedenmalm, H., Makarov, N., *Fluctuations of eigenvalues of random normal matrices*, Preprint in 2008 at arXiv.org/abs/math.PR/0807375. (Now submitted.)
- [2] Ameur, Y., Hedenmalm, H., Makarov, N., *Random normal matrices and Ward's identities*, To appear.
- [3] Berman, R., *Bergman kernels and weighted equilibrium measures of $\mathbb{C}^n$* . Indiana Univ. J. Math. **5** (2009).
- [4] Berman, R., *Determinantal point processes and fermions on complex manifolds: bulk universality*, Preprint in 2008 at arXiv.org/abs/math.CV/08113341.
- [5] Berman, R., *Super Toeplitz operators on holomorphic line bundles*, J. Geom. Anal. **16** (2006), 1–22.
- [6] Berman, R., Berndtsson, B., Sjöstrand, J., *A direct approach to Bergman kernel asymptotics for positive line bundles*, Ark. Mat. **46** (2008).
- [7] Berndtsson, B., *Uniform estimates with weights for the $\bar{\partial}$ -equation*, J. Geom. Anal. **7**, 195–215 (1997).
- [8] Berndtsson, B., *Weighted estimates for the $\bar{\partial}$ -equation*, Complex Analysis and Geometry, 43–57, Ohio State Univ. Math. Res. Publ. **9**, de Gruyter, Berlin, 2001.
- [9] Boutet de Monvel, L., Sjöstrand, J., *Sur la singularité des noyaux de Bergman et de Szegő*, Astérisque **34-35** (1976) p. 123–164.
- [10] Berezin, F. A., *Quantization*, Math. USSR Izvostiya **8** (1974), 1109–1163.
- [11] Duren, P., Schuster, A., *Bergman spaces*, Mathematical surveys and monographs **100**, 2004.
- [12] Engliš, M., *Weighted Bergman kernels and quantization*, Commun. Math. Phys. **227**, 211–241 (2002).
- [13] Fefferman, C., *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*, Invent. Math. **26**, 1–65 (1974).
- [14] Garnett, J. B., Marshall, D. E., *Harmonic measure*, Cambridge 2005.
- [15] Hedenmalm, H., Korenblum, B., Zhu, K., *Theory of Bergman spaces*, Springer 2000.
- [16] Hedenmalm, H., Makarov, N., *Quantum Hele-Shaw flow*, Preprint in 2004 at arXiv.org/abs/math.PR/0411437.
- [17] Hörmander, L., *An introduction to complex analysis in several variables*, Third edition, North-Holland 1990.
- [18] Hörmander, L., *Notions of convexity*, Birkhäuser 1994.
- [19] Jakobsson, S., *Singularity resolution of weighted Bergman kernels*, Contemp. Math. **404**, 121–144.
- [20] Lindholm, L., *Sampling in weighted L^p -spaces of entire functions in $\mathbb{C}^n$ and estimates of the Bergman kernel*, J. Funct. Anal. **182** (2001), 390–426.
- [21] Saff, E. B., Totik, V., *Logarithmic potentials with external fields*, Springer 1997.
- [22] Sjöstrand, J., *Singularités analytiques microlocales*, Astérisque **95** (1982), 1–166.
- [23] Szegő, G., *Über eine Eigenschaft der Exponentialreihe*, Sitzungber. Berlin Math. Gesellschaftswiss. **23** (1924), 50–64.

YACIN AMEUR, DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY,, Box 480, 751 06 UPPSALA, SWEDEN
E-mail address: yacin.ameur@gmail.com

HÅKAN HEDENMALM, DEPARTMENT OF MATHEMATICS, THE ROYAL INSTITUTE OF TECHNOLOGY, S – 100 44 STOCKHOLM, SWEDEN
E-mail address: haakanh@math.kth.se

NIKOLAI MAKAROV, DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125, USA
E-mail address: makarov@caltech.edu